arXiv:1011.5759v3 [math.RT] 31 Aug 2012

QUIVER VARIETIES AND ADJOINT CRYSTALS OF LEVEL ℓ FOR TYPE $A_n^{(1)}$

EUIYONG PARK [1]

Abstract. Let $B(\Lambda)$ be a level ℓ highest weight crystal of the quantum affine algebra $U_q(A_n^{(1)})$. We construct an explicit crystal isomorphism between the geometric realization $\mathfrak{B}(\Lambda)$ of the crystal $B(\Lambda)$ using quiver varieties and the path realization $\mathcal{P}^{\mathrm{ad}}(\Lambda)$ arising from the adjoint crystal $B^{\mathrm{ad},\ell}$ of level ℓ.

Introduction

The *theory of perfect crystals* was developed in studying vertex models in terms of representation theory of quantum affine algebras [10]. The theory of perfect crystals has a lot of applications to several research areas, and plays an important role in the theory of crystal bases. The crystal $B(\Lambda)$ of an irreducible highest weight module $V(\Lambda)$ with highest weight Λ over a quantum affine algebra can be realized as the crystal $\mathcal{P}^B(\Lambda)$ consisting of Λ-paths in a perfect crystal B. In [1], Benkart, Frenkel, Kang and Lee gave a uniform construction of level 1 perfect crystals $B^{\mathrm{ad},1} = B(0) + B(\theta)$, called the *adjoint crystals*, for all quantum affine algebras. Here, θ is the maximal (short) root of the corresponding finite-dimensional simple Lie algebras. It was expected that the crystal $B^{\mathrm{ad},\ell} = B(0) + B(\theta) + \cdots + B(\ell\theta)$ is a perfect crystal of level ℓ. This conjecture was proved for types $A_n^{(1)}, C_n^{(1)}, A_{2n}^{(2)}, D_{n+1}^{(2)}$ and $D_n^{(1)}$ (which yield types $B_n^{(1)}, A_{2n-1}^{(2)}$) in [9, 11, 18, 26]. These perfect crystals are called the *adjoint crystals of level ℓ*.

Let $\mathfrak{g}$ be a symmetric Kac-Moody algebra. Lusztig [19, 20] gave a geometric construction of $U_q^-(\mathfrak{g})$ in terms of perverse sheaves and introduced the notion of *canonical bases* (or *lower global bases* by Kashiwara [15]). In [17], Kashiwara and Saito proved that the set $\mathfrak{B}(\infty)$ of irreducible components of Lusztig's quiver varieties has a crystal structure, which is isomorphic to the crystal $B(\infty)$ of $U_q^-(\mathfrak{g})$. For a dominant integral weight Λ, Nakajima [22, 23] defined a new family of quiver varieties corresponding to Λ and constructed a geometric realization of $V(\Lambda)$. In [24], Saito defined a crystal structure on the set $\mathfrak{B}(\Lambda)$ of irreducible components of certain Lagrangian subvarieties of Nakajima's quiver varieties, and showed that $\mathfrak{B}(\Lambda)$ is isomorphic to the crystal $B(\Lambda)$.

In this paper, we give an explicit crystal isomorphism from the geometric realization $\mathfrak{B}(\Lambda)$ to the path realization $\mathcal{P}^{\mathrm{ad}}(\Lambda)$ arising from the adjoint crystal $B^{\mathrm{ad},\ell}$ of level ℓ for type $A_n^{(1)}$. This is a generalization of the previous result [13]. Let $\mathfrak{g}$ be the Kac-Moody algebra of type $A_n^{(1)}$ and $\mathring{\mathfrak{g}}$ be the

---

2000 *Mathematics Subject Classification.* 17B10, 17B67, 81R10.

Key words and phrases. quiver varieties, adjoint crystals, crystal graphs.

[1] This work was supported by KRF Grant # 2007-341-C00001.

corresponding simple Lie algebra. Let $B^{1,\ell}$ (resp. $B^{n,\ell}$) be the perfect crystal of level ℓ given by (2.1) (resp. (2.4)). Choose the counterclockwise orientation Ω of the Dynkin diagram of type $A_n^{(1)}$. Let Λ be a dominant integral weight of level ℓ, and let $\Lambda' := \sum_{i\in I}\Lambda(h_{i+1})\Lambda_i$ and $\Lambda'' := \sum_{i\in I}\Lambda(h_{i-1})\Lambda_i$, where Λ_i is the ith fundamental weight. We first deduce two kinds of *Young walls* [5, 8] from the results in [3, 7, 12], and describe irreducible components of quiver varieties by using these two kinds of Young walls; see Proposition 4.2, Proposition 4.3. Then, using these descriptions, we give a geometric interpretation of the *fundamental isomorphism of perfect crystals*

$$\mathfrak{B}(\Lambda) \simeq \mathfrak{B}(\Lambda') \otimes B^{1,\ell} \ \text{ and } \ \mathfrak{B}(\Lambda) \simeq \mathfrak{B}(\Lambda'') \otimes B^{n,\ell}$$

(Theorem 5.4). Unlike the approach in [13], we deal directly with irreducible components in the crystal $\mathfrak{B}(\Lambda)$ without using the strict embedding $\mathfrak{B}(\Lambda) \hookrightarrow \mathfrak{B}(\infty) \otimes T_\Lambda \otimes C$. From these fundamental isomorphisms, we give explicit isomorphisms from $\mathfrak{B}(\Lambda)$ to the path realizations $\mathcal{P}^1(\Lambda)$ and $\mathcal{P}^n(\Lambda)$ arising from $B^{1,\ell}$ and $B^{n,\ell}$ in terms of dimension vectors (Theorem 5.6). Using the crystal isomorphism $B^{\mathrm{ad},\ell} \simeq B^{1,\ell} \otimes B^{n,\ell}$, we obtain a geometric interpretation of the isomorphism

$$\mathfrak{B}(\Lambda) \simeq \mathfrak{B}(\Lambda) \otimes B^{\mathrm{ad},\ell}$$

and an explicit map

$$\mathfrak{B}(\Lambda) \xrightarrow{\sim} \mathcal{P}^{\mathrm{ad}}(\Lambda)$$

in terms of dimension vectors; see Theorem 5.7 and Corollary 5.8.

This paper is organized as follows. Section 1 contains the notion of crystal for $U_q(A_n^{(1)})$ and the theory of perfect crystals. Section 2 gives a description of the adjoint crystals $B^{\mathrm{ad},\ell}$ of level ℓ for type $A_n^{(1)}$ in terms of *Young tableaux.* In Section 3, we review the geometric realizations $\mathfrak{B}(\infty)$ and $\mathfrak{B}(\Lambda)$ using quiver varieties. In Section 4, we deduce two kinds of Young walls $\mathcal{Y}^1(\Lambda)$ and $\mathcal{Y}^n(\Lambda)$ which give combinatorial realizations of $B(\Lambda)$, and investigate connections between $\mathcal{Y}^1(\Lambda)$ (resp. $\mathcal{Y}^n(\Lambda)$) and $\mathfrak{B}(\Lambda)$. Then we describe irreducible components of quiver varieties by using these two kinds of Young walls. In Section 5, we construct an explicit crystal isomorphism $\mathfrak{B}(\Lambda) \xrightarrow{\sim} \mathcal{P}^{\mathrm{ad}}(\Lambda)$. We first find geometric interpretations of the isomorphisms $\mathfrak{B}(\Lambda) \simeq \mathfrak{B}(\Lambda') \otimes B^{1,\ell}$ and $\mathfrak{B}(\Lambda) \simeq \mathfrak{B}(\Lambda'') \otimes B^{n,\ell}$ from the description using Young walls in Section 4. Then we give an explicit description of the Λ-path in $B^{1,\ell}$ (resp. $B^{n,\ell}$) corresponding to an irreducible component $X \in \mathfrak{B}(\Lambda)$ in terms of dimension vectors. Using the crystal isomorphism $B^{\mathrm{ad},\ell} \simeq B^{1,\ell} \otimes B^{n,\ell}$, we obtain an explicit isomorphism $\mathfrak{B}(\Lambda) \xrightarrow{\sim} \mathcal{P}^{\mathrm{ad}}(\Lambda)$ in terms of dimension vectors.

Acknowledgments. The author would like to express his deepest gratitude to Professor Seok-Jin Kang for his guidance, valuable comments and enthusiastic encouragement.

1. CRYSTAL GRAPHS FOR $U_q(A_n^{(1)})$

Let $I := \mathbb{Z}/(n+1)\mathbb{Z}$ be an index set. The *affine Cartan datum of type* $A_n^{(1)}$ consists of (i) the *affine Cartan matrix* $A = (a_{ij})_{i,j\in I}$ of type $A_n^{(1)}$, (ii) *dual weight lattice* $P^\vee := \bigoplus_{i=0}^n \mathbb{Z}h_i \oplus \mathbb{Z}d$, (iii) *affine weight lattice* $P := \bigoplus_{i=0}^n \mathbb{Z}\Lambda_i \oplus \mathbb{Z}\delta \subset \mathfrak{h}^*$, where $\mathfrak{h} := \mathbb{C} \otimes_{\mathbb{Z}} P^\vee$, $\Lambda_i(h_j) = \delta_{ij}$, $\Lambda_i(d) = 0$, $\delta(h_i) =$

0, $\delta(d) = 1$ $(i, j \in I)$, (iv) the set of *simple coroots* $\Pi^\vee = \{h_i |\ i \in I\}$, (v) the set of *simple roots* $\Pi = \{\alpha_i |\ i \in I\}$ given by

$$\alpha_j(h_i) = a_{ij} \ \text{ and } \alpha_j(d) = \delta_{0,j} \ (i, j \in I).$$

The free abelian group $Q := \bigoplus_{i=0}^n \mathbb{Z}\alpha_i$ is called the *root lattice* and the semigroup $Q^+ := \sum_{i=0}^n \mathbb{Z}_{\geq 0}\alpha_i$ is called the *positive root lattice*. For $\alpha = \sum_{i\in I} k_i\alpha_i \in Q^+$, the number $\mathrm{ht}(\alpha) := \sum_{i\in I} k_i$ is called the *height* of α. The elements in $P^+ := \{\lambda \in P \mid \lambda(h_i) \geq 0,\ i \in I\}$ are called the *dominant integral weights.*

Let $U_q(\mathfrak{g})$ be the *quantum affine algebra* associated with the affine Cartan datum $(A, \Pi, \Pi^\vee, P, P^\vee)$ of type $A_n^{(1)}$. Let $U_q^-(\mathfrak{g})$ (resp. $U_q^+(\mathfrak{g})$) be the subalgebra of $U_q(\mathfrak{g})$ generated by f_i (resp. e_i) for $i \in I$.

Definition 1.1. A *crystal* B associated with $U_q(\mathfrak{g})$ is a set together with the maps $\mathrm{wt} : B \to P$, $\tilde{e}_i, \tilde{f}_i : B \to B \sqcup \{0\}$, and $\varepsilon_i, \varphi_i : B \to \mathbb{Z} \cup \{-\infty\}$ $(i \in I)$ satisfying the following conditions:

(1) $\varphi_i(b) = \varepsilon_i(b) + \langle h_i, \mathrm{wt}(b)\rangle$ for all $i \in I$,
(2) $\mathrm{wt}(\tilde{e}_i b) = \mathrm{wt}(b) + \alpha_i$ if $\tilde{e}_i b \in B$,
(3) $\mathrm{wt}(\tilde{f}_i b) = \mathrm{wt}(b) - \alpha_i$ if $\tilde{f}_i b \in B$,
(4) $\varepsilon_i(\tilde{e}_i b) = \varepsilon_i(b) - 1,\ \varphi_i(\tilde{e}_i b) = \varphi_i(b) + 1$ if $\tilde{e}_i b \in B$,
(5) $\varepsilon_i(\tilde{f}_i b) = \varepsilon_i(b) + 1,\ \varphi_i(\tilde{f}_i b) = \varphi_i(b) - 1$ if $\tilde{f}_i b \in B$,
(6) $\tilde{f}_i b = b'$ if and only if $b = \tilde{e}_i b'$ for $b, b' \in B,\ i \in I$,
(7) if $\varphi_i(b) = -\infty$ for $b \in B$, then $\tilde{e}_i b = \tilde{f}_i b = 0$.

We refer the reader to [5, 14] for the *tensor product rule* of crystals. For $\Lambda \in P^+$, let $V(\Lambda)$ be the irreducible highest weight $U_q(\mathfrak{g})$-module with highest weight Λ. It was shown in [15] that $U_q^-(\mathfrak{g})$ and $V(\Lambda)$ have crystal bases, denoted by $B(\infty)$ and $B(\Lambda)$ respectively.

We now recall the notion of perfect crystals. Let $U_q'(\mathfrak{g})$ be the subalgebra of $U_q(\mathfrak{g})$ generated by $e_i, f_i, q^{\pm h_i}$ $(i \in I)$, and set $\overline{P} := \bigoplus_{i=0}^n \mathbb{Z}\Lambda_i$, $\overline{P}^+ := \sum_{i=0}^n \mathbb{Z}_{\geq 0}\Lambda_i$, $\overline{P}^\vee := \bigoplus_{i=0}^n \mathbb{Z}h_i$ and $\overline{\mathfrak{h}} := \mathbb{C} \otimes_{\mathbb{Z}} \overline{P}^\vee$. Denote by $\mathrm{cl} : P \to \overline{P}$ the natural projection from P to $\overline{P}$. Given a $U_q'(\mathfrak{g})$-crystal B and $b \in B$, let

$$\varepsilon(b) := \sum_{i=0}^n \varepsilon_i(b)\Lambda_i, \qquad \varphi(b) := \sum_{i=0}^n \varphi_i(b)\Lambda_i.$$

Definition 1.2. A *perfect crystal of level* ℓ is a finite $U_q'(\mathfrak{g})$-crystal B satisfying the following conditions:

(1) there exists a finite dimensional $U_q'(\mathfrak{g})$-module with a crystal basis whose crystal graph is isomorphic to B,
(2) $B \otimes B$ is connected,
(3) there exists a classical weight $\Lambda' \in \overline{P}$ such that

$$\mathrm{wt}(B) \subset \Lambda' + \sum_{i\neq 0} \mathbb{Z}_{\leq 0}\alpha_i, \qquad \#(B_{\Lambda'} := \{b \in B \mid \mathrm{wt}(b) = \Lambda'\}) = 1,$$

(4) for any $b \in B$, we have $\langle c := \sum_{i\in I} h_i,\ \varepsilon(b)\rangle \geq \ell$,
(5) for each $\Lambda \in \overline{P}_\ell^+ := \{\mu \in \overline{P}^+ |\ \langle c,\ \mu\rangle = \ell\}$, there exist unique vectors b^Λ and b_Λ in B such that $\varepsilon(b^\Lambda) = \Lambda,\ \varphi(b_\Lambda) = \Lambda$.

Given $\Lambda \in P^+$ with level ℓ and a perfect crystal B of level ℓ, it was shown in [10, 11] that there exists a unique crystal isomorphism, called the *fundamental isomorphism of perfect crystals*,

$$\Phi : B(\Lambda) \xrightarrow{\sim} B(\varepsilon(b_\Lambda)) \otimes B \tag{1.1}$$

sending the highest weight vector u_Λ to $u_{\varepsilon(b_\Lambda)} \otimes b_\Lambda$. By applying this crystal isomorphism repeatedly, we get a sequence of crystal isomorphisms

$$B(\Lambda) \xrightarrow{\sim} B(\Lambda^1) \otimes B \xrightarrow{\sim} B(\Lambda^2) \otimes B \otimes B \xrightarrow{\sim} \cdots\cdots ,$$

where $\Lambda^0 = \Lambda$, $b_0 = b_\Lambda$, $\Lambda^{k+1} = \varepsilon(b_k)$, $b_{k+1} = b_{\Lambda^{k+1}}$ $(k \geq 0)$. The sequence $\mathbf{p}_\Lambda := (b_k)_{k=0}^\infty$ is called the *ground-state path of weight* Λ and a sequence $\mathbf{p} = (p_k)_{k=0}^\infty$ of elements $p_k \in B$ is called a Λ*-path in* B if $p_k = b_k$ for all $k \gg 0$. Let $\mathcal{P}^B(\Lambda)$ be the set of Λ-paths in B. Then we have the *path realization* of $B(\Lambda)$ arising from B.

Theorem 1.3. [10] *There exists a unique crystal isomorphism* $B(\Lambda) \xrightarrow{\sim} \mathcal{P}^B(\Lambda)$ *which maps* u_Λ *to* $\mathbf{p}_\Lambda$.

2. Adjoint crystals of level ℓ for type $A_n^{(1)}$

In this section, we give a description of adjoint crystals $B^{\mathrm{ad},\ell}$ for type $A_n^{(1)}$ in terms of *Young tableaux* [18]. Let $U_q(\mathring{\mathfrak{g}})$ be the subalgebra of $U_q(\mathfrak{g})$ generated by f_i, e_i and q^{h_i} for $i \in I \setminus \{0\}$, and let $\varpi_i := \Lambda_i - \Lambda_0$ $(1 \leq i \leq n)$. It is well-known that a $U_q(\mathring{\mathfrak{g}})$-crystal can be realized as the set of semistandard tableaux with entries $1, \ldots, n+1$ ([5, Chapter 7], [16], etc).

Let us first recall the perfect crystals $B^{1,\ell}$ and $B^{n,\ell}$ given in [11]. Note that, though $B^{1,\ell}$ and $B^{n,\ell}$ are dual to each other, it is difficult to describe an explicit crystal isomorphism between the path realizations arising from $B^{1,\ell}$ and $B^{n,\ell}$. As a $U_q(\mathring{\mathfrak{g}})$-crystal, $B^{1,\ell}$ is isomorphic to the crystal $B(\ell\varpi_1)$. For $b \in B(\ell\varpi_1)$, let $\nu_i(b)$ be the number of i in b, and set $\nu_0(b) := \nu_{n+1}(b)$. Then

$$B^{1,\ell} = \left\{ b := \begin{array}{|c|c|c|c|c|c|c|}\hline 1 & \cdots & 1 & \cdots & n+1 & \cdots & n+1 \\ \hline \end{array} \;\middle|\; \sum_{i=1}^{n+1} \nu_i(b) = \ell \right\}. \tag{2.1}$$

The crystal structure of $B^{1,\ell}$ is given in [11, Section 1.2]. The map $\mathrm{wt} : B^{1,\ell} \to \mathrm{wt}(B^{1,\ell})$ is bijective. More precisely, let

$$\psi^{1,\ell} : \mathrm{wt}(B^{1,\ell}) \longrightarrow B^{1,\ell} \tag{2.2}$$

be the map sending a weight $\sum_{k \in I} a_k \Lambda_k$ to the tableau b given by

$$\nu_i(b) = \frac{1}{n+1}\left(\ell - \sum_{k=1}^{n} k a_k\right) + \sum_{k=i}^{n} a_k \quad \text{for } i = 1, \ldots, n+1.$$

Then $\psi^{1,\ell}$ is the inverse of $\mathrm{wt} : B^{1,\ell} \to \mathrm{wt}(B^{1,\ell})$. For each $\Lambda = \sum_{k \in I} a_k \Lambda_k \in P^+$ of level ℓ, the ground-state path $\mathbf{p}^1(\Lambda)$ of Λ is given as follows:

$$\mathbf{p}^1(\Lambda) = (b_k^\Lambda)_{k \geq 0}, \tag{2.3}$$

where b_k^{Λ} is the tableau in $B^{1,\ell}$ obtained by $\nu_j(b_k^{\Lambda}) = a_{j+k \mod n}$ for $j = 1, \dots, n+1$. We denote by $\mathcal{P}^1(\Lambda)$ the path realization arising from $B^{1,\ell}$.

We now consider the crystal $B^{n,\ell}$. As a $U_q(\mathring{\mathfrak{g}})$-crystal, $B^{n,\ell}$ is isomorphic to the crystal $B(\ell\varpi_n)$. Let

$$\boxed{\overline{i}} := \begin{array}{|c|}\hline 1 \\ \hline \vdots \\ \hline i-1 \\ \hline i+1 \\ \hline \vdots \\ \hline n+1 \\ \hline \end{array}.$$

For $\overline{b} \in B(\ell\varpi_n)$, let $\overline{\nu}_i(\overline{b})$ denote the number of $\overline{i}$ in $\overline{b}$, and set $\overline{\nu}_0(\overline{b}) := \overline{\nu}_{n+1}(\overline{b})$. Then

$$B^{n,\ell} = \left\{ \overline{b} := \begin{array}{|c|c|c|c|c|c|c|}\hline \overline{n+1} & \cdots & \overline{n+1} & \cdots & \overline{1} & \cdots & \overline{1} \\ \hline \end{array} \;\Big|\; \sum_{i=1}^{n+1} \overline{\nu}_i(\overline{b}) = \ell \right\}. \tag{2.4}$$

The crystal structure of $B^{n,\ell}$ is given in [11, Section 1.2]. Then the map $\mathrm{wt} : B^{n,\ell} \to \mathrm{wt}(B^{n,\ell})$ is bijective, and the inverse is given by

$$\psi^{n,\ell} : \mathrm{wt}(B^{n,\ell}) \longrightarrow B^{n,\ell}, \tag{2.5}$$

which maps a weight $\sum_{k \in I} a_k \Lambda_k$ to the tableau $\overline{b}$ obtained by

$$\overline{\nu}_i(\overline{b}) = \frac{1}{n+1}\left(\ell + \sum_{k=1}^{n} k a_k\right) - \sum_{k=i}^{n} a_k \quad \text{for } i = 1, \dots, n+1.$$

For $\Lambda = \sum_{k \in I} a_k \Lambda_k \in P^+$ of level ℓ, the ground-state path $\mathbf{p}^n(\Lambda)$ of Λ is given as follows:

$$\mathbf{p}^n(\Lambda) = (\overline{b}_k^{\Lambda})_{k \geq 0}, \tag{2.6}$$

where $\overline{b}_k^{\Lambda}$ is the tableau in $B^{n,\ell}$ obtained by $\overline{\nu}_j(\overline{b}_k^{\Lambda}) = a_{j-k-1 \mod n}$ for $j = 1, \dots, n+1$. We denote by $\mathcal{P}^n(\Lambda)$ the path realization arising from $B^{n,\ell}$.

Let $\theta := \varpi_1 + \varpi_n$. Then any tableau T of $B(k\theta)$ $(1 \leq k \leq \ell)$ can be written as

$$T = \begin{array}{|c|c|c|c|c|c|}\hline \overline{i}_1 & \cdots & \overline{i}_k & j_1 & \cdots & j_k \\ \hline \end{array} \tag{2.7}$$

for some $i_1 \geq \cdots \geq i_k$ and $j_1 \leq \cdots \leq j_k$. As a $U_q(\mathring{\mathfrak{g}})$-crystal, let

$$B^{\mathrm{ad},\ell} := \bigoplus_{k=0}^{\ell} B(k\theta).$$

To define the Kashiwara operators $\tilde{e}_0, \tilde{f}_0$ on $B^{\mathrm{ad},\ell}$, we need some notation. For a tableau T and $s \in \mathbb{Z}_{\geq 0}$, let $\mathfrak{R}_s(T)$ (resp. $\mathfrak{L}_s(T)$) be the tableau obtained from T by removing s columns from the left of T (resp. the right of T). In particular, if we take a tableau T of the form (2.7), then we regard $\mathfrak{R}_k(T) = \begin{array}{|c|c|c|}\hline j_1 & \cdots & j_k \\ \hline \end{array}$ (resp. $\mathfrak{L}_k(T) = \begin{array}{|c|c|c|}\hline \overline{i}_1 & \cdots & \overline{i}_k \\ \hline \end{array}$) as an element in the perfect crystal $B^{1,k}$ (resp.

$B^{n,k}$). Given two tableaux $T \in B(k\varpi_i)$ and $T' \in B(k'\varpi_{i'})$, if it exists, $T * T'$ denotes the tableau in $B(k\varpi_i + k'\varpi_{i'})$ obtained from T by adding T' to the right of T. Fix $b \in B(k\theta) \subset B^{\mathrm{ad},\ell}$. Let

$$b_L := \mathfrak{L}_k(b),\ b_R := \mathfrak{R}_k(b),\ \varepsilon^1 := \nu_1(b_R),\ \varphi^1 := \nu_{n+1}(b_R),\ \varepsilon^2 := \overline{\nu}_{n+1}(b_L),\ \varphi^2 := \overline{\nu}_1(b_L).$$

The Kashiwara operators $\tilde{e}_0, \tilde{f}_0$ are given as follows:

$$\tilde{e}_0(b) := \begin{cases} b_L * \tilde{e}_0(b_R) & \text{if } \varphi^1 \geq \varepsilon^2 \text{ and } \varepsilon^1 > 0, \\ (b_L * \boxed{\overline{1}}\,) * (b_R * \boxed{n+1}\,) & \text{if } \varphi^1 \geq \varepsilon^2, \varepsilon^1 = 0 \text{ and } k < \ell \\ \mathfrak{L}_1(\tilde{e}_0(b_L)) * \mathfrak{R}_1(b_R) & \text{if } \varphi^1 < \varepsilon^2 \text{ and } \varepsilon^1 > 0, \\ \tilde{e}_0(b_L) * b_R & \text{if } \varphi^1 < \varepsilon^2 \text{ and } \varepsilon^1 = 0, \\ 0 & \text{otherwise,} \end{cases}$$

$$\tilde{f}_0(b) := \begin{cases} \mathfrak{L}_1(b_L) * \mathfrak{R}_1(\tilde{f}_0(b_R)) & \text{if } \varphi^1 > \varepsilon^2 \text{ and } \varphi^2 > 0, \\ b_L * \tilde{f}_0(b_R) & \text{if } \varphi^1 > \varepsilon^2 \text{ and } \varphi^2 = 0, \\ \tilde{f}_0(b_L) * b_R & \text{if } \varphi^1 \leq \varepsilon^2 \text{ and } \varphi^2 > 0, \\ (\,\boxed{\overline{n+1}} * b_L) * (\,\boxed{1} * b_R) & \text{if } \varphi^1 \leq \varepsilon^2, \varphi^2 = 0 \text{ and } k < \ell, \\ 0 & \text{otherwise.} \end{cases}$$

Then the crystal $B^{\mathrm{ad},\ell}$ with $\tilde{e}_0, \tilde{f}_0$ becomes a $U_q'(\mathfrak{g})$-crystal.

Lemma 2.1. [18, Section 3.3] *Let* $\xi^\ell : B^{1,\ell} \otimes B^{n,\ell} \to B^{\mathrm{ad},\ell}$ *be the map defined by*

$$\xi^\ell(b \otimes \overline{b}) := \begin{cases} \mathfrak{L}_{\overline{\nu}_1(\overline{b})}(\overline{b}) * \mathfrak{R}_{\overline{\nu}_1(\overline{b})}(b) & \text{if } \nu_1(b) \geq \overline{\nu}_1(\overline{b}). \\ \mathfrak{L}_{\nu_1(b)}(\overline{b}) * \mathfrak{R}_{\nu_1(b)}(b) & \text{otherwise,} \end{cases}$$

for $b \in B^{1,\ell}$ *and* $\overline{b} \in B^{n,\ell}$. *Then* ξ^ℓ *is a crystal isomorphism.*

Note that $B^{\mathrm{ad},\ell}$ is perfect by [10, Lemma 4.6.2].

Lemma 2.2. *Let* $\Lambda := \sum_{k\in I} a_k\Lambda_k$ *be a dominant integral weight of level* ℓ. *We set* $\mathtt{b}$ *to be the tableau in* $B((\ell - a_0)\theta)$ *such that the number of* i *in* $\mathtt{b}$ *and the number of* $\overline{i}$ *in* $\mathtt{b}$ *are equal to* a_{i-1} *for* $2 \leq i \leq n+1$. *Then*

$$\mathbf{p}^{\mathrm{ad}}(\Lambda) := \cdots \otimes \mathtt{b} \otimes \mathtt{b} \otimes \mathtt{b}$$

is the ground-state path of Λ *in* $B^{\mathrm{ad},\ell}$.

Proof. We write $\Lambda = \Lambda_{j_1} + \cdots + \Lambda_{j_\ell}$ for $j_1 \leq \ldots \leq j_\ell$. Define a tableau

$$\mathtt{b} := \begin{array}{|c|c|c|c|c|c|}\hline \overline{j_\ell + 1} & \cdots & \overline{j_{a_0+1} + 1} & j_{a_0+1} + 1 & \cdots & j_\ell + 1 \\ \hline \end{array}\,.$$

By construction, $\mathtt{b}$ is contained in $B((\ell - a_0)\theta)$ and such that the number of i in $\mathtt{b}$ and the number of $\overline{i}$ in $\mathtt{b}$ are equal to a_{i-1} for $1 \leq i \leq n+1$. Therefore, the assertion follows from $\varepsilon(\mathtt{b}) = \varphi(\mathtt{b}) = \Lambda$. $\square$

We denote by $\mathcal{P}^{\mathrm{ad}}(\Lambda)$ the path realization arising from $B^{\mathrm{ad},\ell}$. From (2.2), (2.5) and Lemma 2.1, we obtain the 1-1 correspondence between $\mathrm{wt}(B^{1,\ell}) \times \mathrm{wt}(B^{n,\ell})$ and $B^{\mathrm{ad},\ell}$:

$$\begin{aligned} \psi^{\mathrm{ad},\ell} : \mathrm{wt}(B^{1,\ell}) \times \mathrm{wt}(B^{n,\ell}) &\longrightarrow B^{\mathrm{ad},\ell}, \\ (\alpha, \beta) \qquad &\longmapsto \xi^\ell(\psi^{1,\ell}(\alpha) \otimes \psi^{n,\ell}(\beta)). \end{aligned} \tag{2.8}$$

Example 2.3. Let $\mathfrak{g}$ be of type $A_2^{(1)}$ and $\Lambda := 2\Lambda_0 + \Lambda_1$. We set $b := \tilde{f}_1^4 \tilde{f}_2^5 \tilde{f}_1^2 \tilde{f}_0^4 \tilde{f}_2 \tilde{f}_1 u_\Lambda$ in $B(\Lambda)$, where u_Λ is the maximal vector of $B(\Lambda)$. Then the corresponding elements of b in $\mathcal{P}^1(\Lambda)$, $\mathcal{P}^n(\Lambda)$ and $\mathcal{P}^{\mathrm{ad}}(\Lambda)$ are given as follows:

$$\mathbf{p}^1 = \cdots \otimes \begin{array}{|c|c|c|}\hline 2 & 2 & 3 \\ \hline \end{array} \otimes \begin{array}{|c|c|c|}\hline 2 & 3 & 3 \\ \hline \end{array} \otimes \begin{array}{|c|c|c|}\hline 2 & 2 & 3 \\ \hline \end{array} \otimes \begin{array}{|c|c|c|}\hline 3 & 3 & 3 \\ \hline \end{array} \otimes \begin{array}{|c|c|c|}\hline 1 & 2 & 3 \\ \hline \end{array}\,,$$

$$\mathbf{p}^n = \cdots \otimes \begin{array}{|c|c|c|}\hline \overline{3} & \overline{3} & \overline{1} \\ \hline \end{array} \otimes \begin{array}{|c|c|c|}\hline \overline{2} & \overline{2} & \overline{2} \\ \hline \end{array} \otimes \begin{array}{|c|c|c|}\hline \overline{1} & \overline{1} & \overline{1} \\ \hline \end{array} \otimes \begin{array}{|c|c|c|}\hline \overline{3} & \overline{3} & \overline{2} \\ \hline \end{array} \otimes \begin{array}{|c|c|c|}\hline \overline{1} & \overline{1} & \overline{1} \\ \hline \end{array} \otimes \begin{array}{|c|c|c|}\hline \overline{2} & \overline{1} & \overline{1} \\ \hline \end{array}\,,$$

$$\mathbf{p}^{\mathrm{ad}} = \cdots \otimes \begin{array}{|c|c|}\hline 1 & 2 \\ \hline 3 & \multicolumn{1}{|c}{} \\ \cline{1-1} \end{array} \otimes \begin{array}{|c|c|}\hline 1 & 2 \\ \hline 3 & \multicolumn{1}{|c}{} \\ \cline{1-1} \end{array} \otimes \begin{array}{|c|c|}\hline 2 & 3 \\ \hline 3 & \multicolumn{1}{|c}{} \\ \cline{1-1} \end{array} \otimes \begin{array}{|c|c|c|c|c|c|}\hline 1 & 2 & 2 & 2 & 2 & 3 \\ \hline 3 & 3 & 3 & \multicolumn{3}{|c}{} \\ \cline{1-3} \end{array}\,.$$

3. Geometric realizations of crystals

In this section, we review the geometric constructions $\mathfrak{B}(\Lambda)$ for type $A_n^{(1)}$. Let $I = \mathbb{Z}/(n+1)\mathbb{Z}$ and H the set of the arrows such that $i \to j$ with $i, j \in I,\ i - j = \pm 1$. For $h \in H$, we denote by $\mathrm{in}(h)$ (resp. $\mathrm{out}(h)$) the incoming (resp. outgoing) vertex of h. Define an involution $^- : H \to H$ to be the map interchanging $i \to j$ and $j \to i$. Let $\Omega := \{h \in H \mid \mathrm{in}(h) - \mathrm{out}(h) = 1\}$ so that $H = \Omega \sqcup \overline{\Omega}$; i.e.,



,



We take the map $\epsilon : H \to \{-1, 1\}$ given by $\epsilon(h) := 1$ if $h \in \Omega$ and $\epsilon(h) := -1$ if $h \in \overline{\Omega}$. Given an I-graded vector space $V = \bigoplus_{i=0}^n V_i$, let $\underline{\dim} V := \sum_{i=0}^n \dim(V_i)\alpha_i$. For $\alpha = \sum_{i=0}^n k_i \alpha_i \in Q^+$, let $V(\alpha) := \bigoplus_{i=0}^n V_i(\alpha)$ be an I-graded vector space such that $V_i(\alpha)$ is a $\mathbb{C}$-vector space with an ordered basis $v^i(\alpha) = \{v_0^i, v_1^i \ldots, v_{k_i-1}^i\}$ for $i \in I$. Fix an ordered basis

$$v(\alpha) = \{v_0^0, \ldots, v_{k_0-1}^0, v_0^1, \ldots, v_{k_1-1}^1, \ldots, v_0^n, \ldots v_{k_n-1}^n\},$$

for $V(\alpha)$. For simplicity, we write V (resp. V_i) for $V(\alpha)$ (resp. $V_i(\alpha)$). Let

$$E(\alpha) := E_\Omega(\alpha) \oplus E_{\overline{\Omega}}(\alpha),$$

where $E_\Omega(\alpha) := \bigoplus_{i \in I} \mathrm{Hom}(V_{i-1},\ V_i)$ and $E_{\overline{\Omega}}(\alpha) := \bigoplus_{i \in I} \mathrm{Hom}(V_i,\ V_{i-1})$.

Let π_Ω (resp. $\pi_{\overline{\Omega}}$) denote the natural projection from $E(\alpha)$ to $E_\Omega(\alpha)$ (resp. $E_{\overline{\Omega}}(\alpha)$). For an element $\chi \in E(\alpha)$, if there is no danger of confusion, we write $x = (x_i \in \mathrm{Hom}(V_{i-1},\ V_i))_{i \in I}$ (resp. $\overline{x} = (\overline{x}_i \in \mathrm{Hom}(V_i,\ V_{i-1}))_{i \in I}$) for $\pi_\Omega(\chi)$ (resp. $\pi_{\overline{\Omega}}(\chi)$). The matrix representations of x and $\overline{x}$ in the ordered basis $v(\alpha)$ are given as

$$x = \begin{pmatrix} 0 & & \cdots & 0 & x_0 \\ x_1 & 0 & & & 0 \\ 0 & x_2 & 0 & & \vdots \\ \vdots & & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & x_n & 0 \end{pmatrix}, \qquad \overline{x} = \begin{pmatrix} 0 & \overline{x}_1 & 0 & \cdots & 0 \\ & 0 & \overline{x}_2 & & \vdots \\ \vdots & & 0 & \ddots & 0 \\ 0 & & & \ddots & \overline{x}_n \\ \overline{x}_0 & 0 & \cdots & 0 & 0 \end{pmatrix},$$

where x_i (resp. $\overline{x}_i$) is the matrix representation of $x|_{V_{i-1}} : V_{i-1} \to V_i$ (resp. $\overline{x}|_{V_i} : V_i \to V_{i-1}$) in the ordered bases $v^{i-1}(\alpha)$ and $v^i(\alpha)$ (resp. $v^i(\alpha)$ and $v^{i-1}(\alpha)$).

The algebraic group $G(\alpha) := \prod_{i\in I} \mathrm{Aut}(V_i) \subset \mathrm{Aut}(V)$ acts on $E(\alpha)$ by $g \cdot \chi := g\chi g^{-1}$ for $g \in G(\alpha)$ and $\chi \in E(\alpha)$. For $\chi \in E(\alpha)$ and $h \in H$, define $\chi_h := x_{\mathrm{in}(h)}$ if $h \in \Omega$ and $\chi_h := x_{\mathrm{out}(h)}$ if $h \in \overline{\Omega}$. Let $\langle\ ,\ \rangle$ be the sympletic form on $E(\alpha)$ defined by $\langle \chi, \chi' \rangle := \sum_{h\in H} \epsilon(h)\mathrm{tr}(\chi_h \chi'_{\overline{h}})$ for $\chi, \chi' \in E(\alpha)$. Note that $E(\alpha)$ may be viewed as the cotangent bundle of $E_\Omega(\alpha)$ (resp. $E_{\overline{\Omega}}(\alpha)$) under this form. The *moment map* $\mu = (\mu_i : E(\alpha) \to \mathrm{End}(V_i))_{i\in I}$ is given by $\mu_i(\chi) := \sum_{h\in H,\ \mathrm{in}(h)=i} \epsilon(h)\chi_h\chi_{\overline{h}}$ for $\chi \in E(\alpha)$. Note that, for $\chi = x + \overline{x} \in E(\alpha)$,

$$\mu_i(\chi) = 0 \text{ for all } i \in I \quad \text{if and only if} \quad x\overline{x} - \overline{x}x = 0. \tag{3.1}$$

An element $\chi \in E(\alpha)$ is *nilpotent* if there exists an $N \geq 2$ such that for any sequence $h_1, \ldots, h_N \in H$ satisfying $\mathrm{in}(h_i) = \mathrm{out}(h_{i+1})$ $(i = 1, \ldots, N-1)$, the composition map $\chi_{h_N} \cdots \chi_{h_1}$ is zero. We define *Lusztig's quiver variety* [19, 20] to be

$$\mathfrak{X}(\alpha) := \{\chi \in E(\alpha) \mid \chi : \text{nilpotent},\ \mu_i(\chi) = 0 \text{ for all } i \in I\}.$$

Let $\mathrm{Irr}\mathfrak{X}(\alpha)$ denote the set of all irreducible components of $\mathfrak{X}(\alpha)$. In [17], Kashiwara and Saito gave a crystal structure to $\mathfrak{B}(\infty) := \bigsqcup_{\alpha\in Q^+} \mathrm{Irr}\mathfrak{X}(\alpha)$, and proved the following theorem.

Theorem 3.1. [17, Theorem 5.3.2] *There is a unique crystal isomorphism* $\mathfrak{B}(\infty) \simeq B(\infty)$.

We now review Nakajima's quiver varieties [22, 23]. For $\Lambda = \sum_{i=0}^n a_i\Lambda_i \in P^+$, let $W(\Lambda) := \bigoplus_{i=0}^n W_i(\Lambda)$ such that $W_i(\Lambda)$ is a $\mathbb{C}$-vector space of dimension a_i. For simplicity, we write W (resp. W_i) for $W(\Lambda)$ (resp. $W_i(\Lambda)$). Let

$$X(\Lambda, \alpha) := \mathfrak{X}(\alpha) \times \sum_{i\in I} \mathrm{Hom}(V_i, W_i).$$

The group $G(\alpha)$ acts on $X(\Lambda, \alpha)$ by $g\cdot(\chi, t) := (g\chi g^{-1}, tg^{-1})$. For $\chi \in \mathfrak{X}(\alpha)$, an I-graded subspace S of $V(\alpha)$ is χ-*invariant* if $\chi_h(S_{\mathrm{out}(h)}) \subset S_{\mathrm{in}(h)}$ for all $h \in H$. An element $(\chi, t) \in X(\Lambda, \alpha)$ is called a *stable point* if it satisfies the following conditions: if S is a χ-invariant subspace of V with $t_i(S_i) = 0$ $(i \in I)$, then $S = 0$. For $(\chi = x + \overline{x}, t) \in X(\Lambda, \alpha)$, one can prove that

$$(\chi, t) \text{ is stable } \iff \ker x \cap \ker \overline{x} \cap \ker t = 0. \tag{3.2}$$

Let $X(\Lambda, \alpha)^{\mathrm{st}}$ be the set of all stable points of $X(\Lambda, \alpha)$. We define

$$\mathfrak{X}(\Lambda, \alpha) := X(\Lambda, \alpha)^{\mathrm{st}}/G(\alpha).$$

Let $\mathrm{Irr}\mathfrak{X}(\Lambda, \alpha)$ (resp. $\mathrm{Irr}X(\Lambda, \alpha)$) be the set of all irreducible components of $\mathfrak{X}(\Lambda, \alpha)$ (resp. $X(\Lambda, \alpha)$). Since $\mathrm{Irr}\mathfrak{X}(\Lambda, \alpha)$ can be identified with $\{Y \in \mathrm{Irr}X(\Lambda, \alpha) \mid Y \cap X(\Lambda, \alpha)^{\mathrm{st}} \neq \emptyset\}$, we have

$$\begin{aligned} \Pi_\Lambda : \mathrm{Irr}\mathfrak{X}(\alpha) &\longrightarrow \mathrm{Irr}\mathfrak{X}(\Lambda, \alpha) \sqcup \{0\}, \\ X_0 &\longmapsto \left(\left(X_0 \times \sum_{i\in I} \mathrm{Hom}(V_i, W_i) \right) \cap X(\Lambda, \alpha)^{\mathrm{st}} \right) / G(\alpha). \end{aligned} \tag{3.3}$$

In [24], Saito defined a crystal structure on $\mathfrak{B}(\Lambda) := \bigsqcup_{\alpha\in Q^+} \mathrm{Irr}\mathfrak{X}(\Lambda, \alpha)$, and proved the following theorem.

Theorem 3.2. [24, Theorem 4.6.4] *There is a unique crystal isomorphism* $\mathfrak{B}(\Lambda) \simeq B(\Lambda)$.

4. Quiver varieties and Young walls

The *Young walls* are new combinatorial objects introduced by Kang [5, 8] for realizing crystals of basic representations for several types. Young walls generalize the notion of colored Young diagrams in [2, 21]. In this section, we deduce two kinds of Young walls from the results given in [3, 7, 12, 25], and describe irreducible components of $\mathfrak{B}(\Lambda)$ by using these two kinds of Young walls. This description will play a crucial role in proving Theorem 5.4.

Let $\mathtt{P}_k^1$ and $\mathtt{P}_k^n$ be the patterns of a fundamental weight Λ_k given as follows:

	$\vdots$	$\vdots$	$\vdots$	$\vdots$		$\vdots$	$\vdots$
	1	2	3	4	$\cdots$	$k+1$	$k+2$
	0	1	2	3	$\cdots$	k	$k+1$
	n	0	1	2	$\cdots$	$k-1$	k

the pattern $\mathtt{P}_k^1$ of weight Λ_k

	$\vdots$	$\vdots$	$\vdots$	$\vdots$		$\vdots$	$\vdots$
	0	n	$n-1$	$n-2$	$\cdots$	$k-1$	$k-2$
	1	0	n	$n-1$	$\cdots$	k	$k-1$
	2	1	0	n	$\cdots$	$k+1$	k

the pattern $\mathtt{P}_k^n$ of weight Λ_k

A Young wall Y in $\mathtt{P}_k^1$ (resp. $\mathtt{P}_k^n$) is a wall consisting of colored blocks stacked by the following rules:

(a) the colored blocks should be stacked in the pattern $\mathtt{P}_k^1$ (resp. $\mathtt{P}_k^n$) of weight Λ_k,

(b) except for the rightmost column, there should be no free space to the right of any block.

We number columns of Y from right to left, denoted by $Y = (y_j)_{j\geq 0}$ where y_j is the jth column of Y. Let $\mathrm{ht}(y_j)$ be the number of blocks in y_j, and $\mathrm{wt}(y_j) := \sum_{i\in I} k_{ij}\alpha_i$ $(j \in \mathbb{Z}_{\geq 0})$, where k_{ij} is the number of i-blocks in the jth column y_j. Let $\mathcal{Y}_k^1$ (resp. $\mathcal{Y}_k^n$) denote the set of all Young walls in the pattern $\mathtt{P}_k^1$ (resp. $\mathtt{P}_k^n$). Fix $\Lambda \in P^+$ of level ℓ. Then Λ can be uniquely written as

$$\Lambda = \Lambda_{i_1} + \cdots + \Lambda_{i_\ell}$$

for some $0 \leq i_1 \leq i_2 \leq \cdots \leq i_\ell \leq n$. For an ℓ-tuple $\mathbf{Y} = (Y^{(1)}, \ldots, Y^{(\ell)})$ of Young walls, let $l_j^{(k)}$ be the length of the jth row of $Y^{(k)}$, and $\mathrm{c}_j^{(k)}$ be the color at the left end of the jth row of $Y^{(k)}$. An ℓ-tuple $\mathbf{Y} = (Y^{(1)}, \ldots, Y^{(\ell)})$ of Young walls is said to be *reduced* if, for all $t > 0$,

$$\{\mathrm{c}_j^{(k)} \mid l_j^{(k)} = t,\ 1 \leq k \leq \ell,\ j \geq 0\} \neq \{0, 1, \ldots, n\}.$$

Let

$$\begin{aligned} \mathcal{Y}^1(\Lambda) := \{\mathbf{Y} = (Y^{(1)}, \ldots, Y^{(\ell)}) \mid Y^{(k)} \in \mathcal{Y}^1_{i_k},\ \mathbf{Y} : \text{reduced}, \\ Y^{(1)} \preceq Y^{(2)} \preceq \cdots \preceq Y^{(\ell)} \preceq Y^{(1)} + n + 1\}. \end{aligned} \tag{4.1}$$

Here, if let $Y^{(k)} = (y_j^{(k)})_{j\geq 0}$, then the order $\preceq$ is given as follows:

$$\begin{aligned} Y^{(k)} \preceq Y^{(k+1)} &\iff \mathrm{ht}(y_j^{(k)}) \leq \mathrm{ht}(y_j^{(k+1)}) + i_{k+1} - i_k \ (j \in \mathbb{Z}_{\geq 0}), \\ Y^{(\ell)} \preceq Y^{(1)} + n + 1 &\iff \mathrm{ht}(y_j^{(\ell)}) \leq \mathrm{ht}(y_j^{(1)}) + i_1 - i_\ell + n + 1 \ (j \in \mathbb{Z}_{\geq 0}). \end{aligned}$$

Then, by [3, Proposition 2.11], we have a 1-1 map

$$\begin{array}{rccl} F^1 : & \mathcal{Y}^1(\Lambda) & \longrightarrow & \mathcal{P}^1(\Lambda), \\ & \mathbf{Y} = (Y^{(1)}, \dots, Y^{(\ell)}) & \longmapsto & \left(\psi^{1,\ell} \left(\mathrm{wt}(b_j^{\Lambda}) - \mathrm{cl} \left(\sum_{k=1}^{\ell} \mathrm{wt}(y_j^{(k)}) \right) \right) \right)_{j \geq 0}, \end{array} \tag{4.2}$$

where $\mathbf{p}^1(\Lambda) = (b_k^{\Lambda})_{k \geq 0}$ is the ground-state path of Λ in $B^{1,\ell}$.

In a similar manner, we let

$$\begin{aligned} \mathcal{Y}^n(\Lambda) := \{ \overline{\mathbf{Y}} = (\overline{Y}^{(1)}, \dots, \overline{Y}^{(\ell)}) \mid \overline{Y}^{(k)} \in \mathcal{Y}^n_{i_k},\ \overline{\mathbf{Y}} : \text{reduced}, \\ \overline{Y}^{(1)} \succeq \overline{Y}^{(2)} \succeq \cdots \succeq \overline{Y}^{(\ell)} \succeq \overline{Y}^{(1)} - n - 1 \}. \end{aligned} \tag{4.3}$$

Here, if let $\overline{Y}^{(k)} = (\overline{y}_j^{(k)})_{j \geq 0}$, then the order $\succeq$ is given as follows:

$$\begin{aligned} \overline{Y}^{(k-1)} \succeq \overline{Y}^{(k)} &\iff \mathrm{ht}(\overline{y}_j^{(k-1)}) \geq \mathrm{ht}(\overline{y}_j^{(k)}) + i_{k-1} - i_k \ (j \in \mathbb{Z}_{\geq 0}), \\ \overline{Y}^{(\ell)} \succeq \overline{Y}^{(1)} - n - 1 &\iff \mathrm{ht}(\overline{y}_j^{(\ell)}) \geq \mathrm{ht}(\overline{y}_j^{(1)}) + i_{\ell} - i_1 - n - 1 \ (j \in \mathbb{Z}_{\geq 0}). \end{aligned}$$

We have a 1-1 map

$$\begin{array}{rccl} F^n : & \mathcal{Y}^n(\Lambda) & \longrightarrow & \mathcal{P}^n(\Lambda), \\ & \overline{\mathbf{Y}} = (\overline{Y}^{(1)}, \dots, \overline{Y}^{(\ell)}) & \longmapsto & \left(\psi^{n,\ell} \left(\mathrm{wt}(\overline{b}_j^{\Lambda}) - \mathrm{cl} \left(\sum_{k=1}^{\ell} \mathrm{wt}(\overline{y}_j^{(k)}) \right) \right) \right)_{j \geq 0}, \end{array} \tag{4.4}$$

where $\mathbf{p}^n(\Lambda) = (\overline{b}_k^{\Lambda})_{k \geq 0}$ is the ground-state path of Λ in $B^{n,\ell}$. Then, $\mathcal{Y}^1(\Lambda)$ and $\mathcal{Y}^n(\Lambda)$ have the crystal structures from $\mathcal{P}^1(\Lambda)$ and $\mathcal{P}^n(\Lambda)$ via the maps F^1 and F^n, respectively.

Proposition 4.1. *For $\Lambda \in P^+$, $\mathcal{Y}^1(\Lambda)$ (resp. $\mathcal{Y}^n(\Lambda)$) is isomorphic to the crystal $B(\Lambda)$.*

Note that the crystal structures of $\mathcal{Y}^1(\Lambda)$ and $\mathcal{Y}^n(\Lambda)$ can be described completely in a combinatorial manner by using [3, Proposition 2.11], [7, Theorem 3.13] and [12, Proposition 6.1]. As a set, $\mathcal{Y}^1(\Lambda)$ is different from $\mathcal{Y}^n(\Lambda)$ even when forgetting colors of Young walls. It is difficult to describe an explicit crystal isomorphism between $\mathcal{Y}^1(\Lambda)$ and $\mathcal{Y}^n(\Lambda)$ in a combinatorial manner [6]. We also remark that the description of $\mathcal{Y}^n(\Lambda)$ is almost the same as that of *Young pyramid* given in [25]. Given $\overline{\mathbf{Y}} = (\overline{Y}^{(1)}, \dots, \overline{Y}^{(\ell)}) \in \mathcal{Y}^n(\Lambda)$, one can obtain the corresponding Young pyramid by flipping $\overline{Y}^{(k)}$ in the diagonal and rearranging them in an appropriate manner.

Now we describe irreducible components of quiver varieties using $\mathcal{Y}^1(\Lambda)$ and $\mathcal{Y}^n(\Lambda)$. Take $\alpha := \sum_{i=0}^{n} k_i \alpha_i \in Q^+$ and write V (resp. V_i) for $V(\alpha)$ (resp. $V_i(\alpha)$). Let $k_{-1} = k_n$ and $k_{n+1} = k_0$. By Proposition 4.1 and Theorem 3.2, there is the unique crystal isomorphism from $\mathcal{Y}^1(\Lambda)$ (resp. $\mathcal{Y}^n(\Lambda)$) to $\mathfrak{B}(\Lambda)$. We will describe these crystal isomorphisms explicitly.

For $s \in I$, we define a linear map $\mathcal{E}_{ij}^s : V_{s-1} \to V_s$ (resp. $\overline{\mathcal{E}}_{ij}^s : V_s \to V_{s-1}$) by

$$\mathcal{E}_{ij}^s(v_k^{s-1}) := \begin{cases} v_j^s & \text{if } k = i, \\ 0 & \text{otherwise,} \end{cases} \qquad \left(\text{resp. } \ \overline{\mathcal{E}}_{ij}^s(v_k^s) := \begin{cases} v_j^{s-1} & \text{if } k = i, \\ 0 & \text{otherwise.} \end{cases} \right)$$

We first deal with an isomorphism of $\mathcal{Y}^n(\Lambda)$ and $\mathfrak{B}(\Lambda)$. Take $\overline{\mathbf{Y}} = (\overline{Y}^{(1)}, \dots, \overline{Y}^{(\ell)}) \in \mathcal{Y}^n(\Lambda)$. Let $\overline{\mathfrak{b}}_{ij}^{(k)}$ denote the ith block from the bottom of the jth column of $\overline{Y}^{(k)}$ and $\mathrm{c}(\overline{\mathfrak{b}}_{ij}^{(k)})$ be the color of $\overline{\mathfrak{b}}_{ij}^{(k)}$. We set $o(\overline{\mathfrak{b}}_{ij}^{(k)}) := \#\{\overline{\mathfrak{b}}_{rs}^{(t)} \in \overline{\mathbf{Y}} \mid \mathrm{c}(\overline{\mathfrak{b}}_{ij}^{(k)}) = \mathrm{c}(\overline{\mathfrak{b}}_{rs}^{(t)}),\ (t, r, s) \prec (k, i, j)\}$, where $\prec$ is the lexicographical order. Define

$$\overline{x}(\overline{\mathbf{Y}}) := \sum_{\overline{\mathfrak{b}}_{ij}^{(k)} \in \overline{\mathbf{Y}}, j>0} \overline{\mathcal{E}}^{\mathrm{c}(\overline{\mathfrak{b}}_{ij}^{(k)})}_{o(\overline{\mathfrak{b}}_{ij}^{(k)}),\ o(\overline{\mathfrak{b}}_{ij-1}^{(k)})} \in E_{\overline{\Omega}}(\alpha). \tag{4.5}$$

Note that, for $t \geq 0$,

$$\ker(\overline{x}(\overline{\mathbf{Y}}))^t = \mathrm{Span}_{\mathbb{C}}\{v^{\mathrm{c}(\overline{\mathfrak{b}}_{ij}^{(k)})}_{o(\overline{\mathfrak{b}}_{ij}^{(k)})} \mid \overline{\mathfrak{b}}_{ij}^{(k)} \in \overline{Y}^{(k)},\ j < t,\ 1 \leq k \leq \ell\}. \tag{4.6}$$

Since $E(\alpha)$ can be viewed as the cotangent bundle of $E_{\overline{\Omega}}(\alpha)$, we let

$$X_0(\overline{\mathbf{Y}}) := \text{the closure of the conormal bundle of the } G(\alpha)\text{-orbit of } \overline{x}(\overline{\mathbf{Y}}). \tag{4.7}$$

Proposition 4.2. *The map* $\overline{\mathbf{Y}} \mapsto \Pi_\Lambda(X_0(\overline{\mathbf{Y}}))$ *from* $\mathcal{Y}^n(\Lambda)$ *to* $\mathfrak{B}(\Lambda)$ *is a crystal isomorphism.*

Proof. By the definition of Young walls, the pair $(\overline{x}(\overline{\mathbf{Y}}), V(\alpha))$ becomes a nilpotent representation of the quiver $(I, \overline{\Omega})$. Moreover, it follows from the definition of $\mathcal{Y}^n(\Lambda)$ in (4.3) that $(\overline{x}(\overline{\mathbf{Y}}), V(\alpha))$ satisfies the *aperiodic condition* given in [20, Section 15.4]. Hence, by [20, Corollary 15.6], $X_0(\overline{\mathbf{Y}})$ is an irreducible component of $\mathfrak{X}(\alpha)$. Therefore, since the description of $\mathcal{Y}^n(\Lambda)$ is almost the same as that of Young pyramid given in [25], our assertion follows from [4, Theorem 6.3] and [25, Theorem 8.4]. □

Now we consider the crystal isomorphism between $\mathcal{Y}^1(\Lambda)$ and $\mathfrak{B}(\Lambda)$. Let $\mathbf{Y} = (Y^{(1)}, \dots, Y^{(\ell)}) \in \mathcal{Y}^1(\Lambda)$. Let $\mathfrak{b}_{ij}^{(k)}$ denote the ith block from the bottom of the jth column of $Y^{(k)}$, and $\mathrm{c}(\mathfrak{b}_{ij}^{(k)})$ be the color of $\mathfrak{b}_{ij}^{(k)}$. We set $o(\mathfrak{b}_{ij}^{(k)}) := \#\{\mathfrak{b}_{rs}^{(t)} \in \mathbf{Y} \mid \mathrm{c}(\mathfrak{b}_{ij}^{(k)}) = \mathrm{c}(\mathfrak{b}_{rs}^{(t)}),\ (t, r, s) \prec (k, i, j)\}$, where $\prec$ is the lexicographical order. Define

$$x(\mathbf{Y}) := \sum_{\mathfrak{b}_{ij}^{(k)} \in \mathbf{Y}, j>0} \mathcal{E}^{\mathrm{c}(\mathfrak{b}_{i,j-1}^{(k)})}_{o(\mathfrak{b}_{ij}^{(k)}),\ o(\mathfrak{b}_{ij-1}^{(k)})} \in E_{\Omega}(\alpha). \tag{4.8}$$

Note that, for $t \geq 0$,

$$\ker(x(\mathbf{Y}))^t = \mathrm{Span}_{\mathbb{C}}\{v^{\mathrm{c}(\mathfrak{b}_{ij}^{(k)})}_{o(\mathfrak{b}_{ij}^{(k)})} \mid \mathfrak{b}_{ij}^{(k)} \in Y^{(k)},\ j < t,\ 1 \leq k \leq \ell\}. \tag{4.9}$$

Since $E(\alpha)$ can be viewed as the cotangent bundle of $E_{\Omega}(\alpha)$, let

$$X_0(\mathbf{Y}) := \text{the closure of the conormal bundle of the } G(\alpha)\text{-orbit of } x(\mathbf{Y}). \tag{4.10}$$

Proposition 4.3. *The map* $\mathbf{Y} \mapsto \Pi_\Lambda(X_0(\mathbf{Y}))$ *from* $\mathcal{Y}^1(\Lambda)$ *to* $\mathfrak{B}(\Lambda)$ *is a crystal isomorphism.*

Proof. Note that $\mathcal{Y}^1(\Lambda)$ and $\mathcal{Y}^n(\Lambda)$ are totally different as sets. By the definition of Young walls, the pair $(x(\mathbf{Y}), V(\alpha))$ becomes a nilpotent representation of the quiver (I, Ω). Since the counterclockwise orientation Ω is opposite to $\overline{\Omega}$, by the symmetry of Ω and $\overline{\Omega}$, one can prove that the pair $(x(\mathbf{Y}), V(\alpha))$ satisfies the aperiodic condition and $X_0(\mathbf{Y})$ is an irreducible component of $\mathfrak{X}(\alpha)$. For the same reason, our assertion can be proved in a similar manner to that of [25, Theorem 8.4]. □

Example 4.4. We keep all notations in Example 2.3. The Young walls corresponding to b in $\mathcal{Y}^1(\Lambda)$ and $\mathcal{Y}^n(\Lambda)$ are given as follows.





We set $\alpha := \Lambda - \mathrm{wt}(b) = 4\alpha_0 + 7\alpha_1 + 6\alpha_2$. Let X be the irreducible component of $\mathfrak{B}(\Lambda)$ corresponding to b via the isomorphism in Theorem 3.2. The matrices $x(\mathbf{Y})$ and $x(\overline{\mathbf{Y}})$ are given as follows.

$$x(\mathbf{Y}) := \mathcal{E}^0_{00} + \mathcal{E}^0_{11} + \mathcal{E}^0_{32} + \mathcal{E}^0_{53} + \mathcal{E}^1_{24} + \mathcal{E}^2_{00} + \mathcal{E}^2_{21} + \mathcal{E}^2_{53} + \mathcal{E}^2_{64},$$
$$\overline{x}(\overline{\mathbf{Y}}) := \overline{\mathcal{E}}^0_{23} + \overline{\mathcal{E}}^1_{00} + \overline{\mathcal{E}}^1_{21} + \overline{\mathcal{E}}^1_{52} + \overline{\mathcal{E}}^1_{63} + \overline{\mathcal{E}}^2_{00} + \overline{\mathcal{E}}^2_{34} + \overline{\mathcal{E}}^2_{45}.$$

Note that $x(\mathbf{Y}) + \overline{x}(\overline{\mathbf{Y}}) \notin \mathfrak{X}(\alpha)$ since $[x(\mathbf{Y}), \overline{x}(\overline{\mathbf{Y}})] \neq 0$.

5. QUIVER VARIETIES AND THE CRYSTALS $B^{1,\ell}$, $B^{n,\ell}$ AND $B^{\mathrm{ad},\ell}$

In this section, we construct an explicit crystal isomorphism from the geometric realization $\mathfrak{B}(\Lambda)$ to the path realization $\mathcal{P}^{\mathrm{ad}}(\Lambda)$ (resp. $\mathcal{P}^1(\Lambda)$, $\mathcal{P}^n(\Lambda)$) arising from $B^{\mathrm{ad},\ell}$ (resp. $B^{1,\ell}$, $B^{n,\ell}$). Let $X \in \mathfrak{B}(\Lambda)$ for $\Lambda \in P^+$ of level ℓ and $\alpha := \Lambda - \mathrm{wt}(X)$. For an element $(\chi, t) \in X(\Lambda, \alpha)$, let $[\chi, t]$ denote the $G(\alpha)$-orbit of (χ, t) in $X(\Lambda, \alpha)$. For an element $[\chi = x + \overline{x}, t]$ in a certain open subset of X, we will give an explicit description of the Λ-path in $B^{\mathrm{ad},\ell}$ (resp. $B^{1,\ell}$, $B^{n,\ell}$) corresponding to X in terms of dimension vectors of $\ker(x\overline{x})^{k+1}/\ker \overline{x}(x\overline{x})^k$ and $\ker \overline{x}(x\overline{x})^k/\ker(x\overline{x})^k$ (resp. $\ker x^{k+1}/\ker x^k$, $\ker \overline{x}^{k+1}/\ker \overline{x}^k$). For this purpose, we need several lemmas. Combining (3.3) with results in [13], we have the following lemmas.

Lemma 5.1. [13, Lemma 5.1] *Let $X \in \mathfrak{B}(\Lambda)$. For any element $[\chi = x + \overline{x}, t] \in X$ and $k \in \mathbb{Z}_{\geq 0}$, we have*

(1) $\ker(x\overline{x})^k = \ker(\overline{x}x)^k$,
(2) $\ker x^k$ *and* $\ker \overline{x}^k$ *are χ-invariant,*
(3) $\ker(x\overline{x})^k$ *is χ-invariant.*

Lemma 5.2. [13, Lemma 5.2] *Let $X \in \mathfrak{B}(\Lambda)$. There is a nonempty open subset $U \subset X$ such that*

$$\ker x^k \simeq \ker x'^k, \quad \ker \overline{x}^k \simeq \ker \overline{x}'^k$$

for any elements $[\chi = x + \overline{x}, t]$, $[\chi' = x' + \overline{x}', t'] \in U$ and $k \in \mathbb{Z}_{\geq 0}$.

Let $M := \bigoplus_{i\in I} M_i$ be an I-graded vector space and $x \in \mathrm{Hom}(M,M)$ be a homogeneous linear map of degree k. For an x-invariant I-graded subspace N of M, let $x|_{M/N}$ (resp. $x|_N$) denote the homogeneous linear map of degree k in $\mathrm{Hom}(M/N, M/N)$ (resp. $\mathrm{Hom}(N,N)$) obtained by restricting x to M/N (resp. N).

Lemma 5.3. *Let $M = \bigoplus_{i\in I} M_i$ and $L = \bigoplus_{i\in I} L_i$. Let $x \in \bigoplus_{i\in I} \mathrm{Hom}(M_{i-1}, M_i)$, and let*

$$N := \ker x \quad \textit{and} \quad x' := x|_{M/N}.$$

Take elements

$$\overline{x}' \in \bigoplus_{i\in I} \mathrm{Hom}(M_i/N_i, M_{i-1}/N_{i-1}) \quad \textit{and} \quad t' \in \bigoplus_{i\in I} \mathrm{Hom}(M_i/N_i, L_{i+1})$$

such that

$$[x', \overline{x}'] = 0 \quad \textit{and} \quad \ker x' \cap \ker \overline{x}' \cap \ker t' = 0,$$

where N_i is the i-subspace of N for $i \in I$. Then there exist

$$\overline{x} \in \bigoplus_{i\in I} \mathrm{Hom}(M_i, M_{i-1}) \quad \textit{and} \quad t \in \bigoplus_{i\in I} \mathrm{Hom}(M_i, L_i)$$

such that

$$[x, \overline{x}] = 0, \quad \overline{x}|_{M/N} = \overline{x}', \quad t' = t \circ x|_{M/N} \quad \textit{and} \quad \ker x \cap \ker \overline{x} \cap \ker t = 0.$$

Proof. By definition, $x|_{\ker x^2/\ker x}$ gives an I-graded isomorphism between $\ker x^2/\ker x$ and $\ker x \cap \mathrm{Im}\ x$. We set

$$\begin{aligned}
\ell &:= \dim L,\\
r &:= \dim(\ker x \cap \mathrm{Im}\ x) = \dim(\ker x^2/\ker x),\\
s &:= \dim(\ker x/\ker x \cap \mathrm{Im}\ x),\\
t &:= \dim(M/\ker x^2).
\end{aligned}$$

Take a homogeneous ordered basis

$$\omega := \{\omega_1^1, \dots, \omega_r^1, \omega_1^2, \dots, \omega_s^2, \omega_1^3, \dots, \omega_r^3, \omega_1^4, \dots, \omega_t^4\}$$

of M such that

$$\begin{aligned}
\ker x \cap \mathrm{Im}\ x &= \mathrm{Span}_{\mathbb{C}}\{\omega_1^1, \dots, \omega_r^1\},\\
\ker x &= \mathrm{Span}_{\mathbb{C}}\{\omega_1^1, \dots, \omega_r^1, \omega_1^2, \dots, \omega_s^2\},\\
\ker x^2 &= \mathrm{Span}_{\mathbb{C}}\{\omega_1^1, \dots, \omega_r^1, \omega_1^2, \dots, \omega_s^2, \omega_1^3, \dots, \omega_r^3\}.
\end{aligned}$$

Then the matrix representative of x with respect to the basis ω is given as follows:

$$x = \begin{pmatrix} 0 & 0 & a & b \\ 0 & 0 & 0 & c \\ 0 & 0 & 0 & d \\ 0 & 0 & 0 & e \end{pmatrix}$$

where a is an $r \times r$ matrix, b, d are $r \times t$ matrices, c is an $s \times t$ matrix and e is a $t \times t$ matrix. Note that a is invertible and $\begin{pmatrix} d \\ e \end{pmatrix}$ has full rank. By construction, the representative of x' is given

$$x' = \begin{pmatrix} 0 & d \\ 0 & e \end{pmatrix}.$$

The matrix representative of $\overline{x}'$ may be written as $\overline{x}' = \begin{pmatrix} f & g \\ 0 & h \end{pmatrix}$. Note that $\ker x^2 / \ker x$ is invariant under $\overline{x}'$ by the assumption $[x', \overline{x}'] = 0$. We have

$$0 = [x', \overline{x}'] = x'\overline{x}' - \overline{x}'x' = \begin{pmatrix} 0 & dh - fd - ge \\ 0 & eh - he \end{pmatrix}. \tag{5.1}$$

Let $X := afa^{-1}$ and Y be an invertible $s \times s$ matrix. Since $\begin{pmatrix} d \\ e \end{pmatrix}$ has full rank, the following equations

$$Z_1 d + Z_3 e = ag + bh - Xb \quad \text{and} \quad Z_2 d + Z_4 e = ch - Yc$$

have solutions. We choose a solution $\begin{pmatrix} Z_1 & Z_3 \\ Z_2 & Z_4 \end{pmatrix}$ of the above equations which maps V_i to V_{i-1} for $i \in I$. We set

$$\overline{x} := \begin{pmatrix} X & 0 & Z_1 & Z_3 \\ 0 & Y & Z_2 & Z_4 \\ 0 & 0 & f & g \\ 0 & 0 & 0 & h \end{pmatrix}$$

By construction, we have $\overline{x}|_{M/\ker x} = \overline{x}'$ and

$$[x, \overline{x}] = \begin{pmatrix} 0 & 0 & af - Xa & ag + bh - Xb - Z_1 d - Z_3 e \\ 0 & 0 & 0 & ch - Yc - Z_2 d - Z_4 e \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} = 0.$$

Now we take a map $t \in \bigoplus_{i \in I} \mathrm{Hom}(M_i, L_i)$ such that, for any element $m = xm' \in \mathrm{Im}\ x$,

$$t(m) = t'(m' + \ker x).$$

Then we have $t \circ x|_{M/\ker x} = t'$. Fix an ordered basis of L, and let $\mathfrak{t}'$ be the matrix representative of $t'|_{\ker x^2/\ker x}$. Since $\ker x' \cap \ker \overline{x}' \cap \ker t' = 0$, we have

$$\ker f \cap \ker \mathfrak{t}' = 0.$$

Then the matrix representative $\mathfrak{t}$ of $t|_{\ker x}$ can be written as

$$\mathfrak{t} = \begin{pmatrix} \mathfrak{t}' a^{-1} & \mathfrak{t}'' \end{pmatrix}$$

for some $r \times \ell$ matrix $\mathfrak{t}''$. Since $\overline{x}|_{\ker x} = \begin{pmatrix} X & 0 \\ 0 & Y \end{pmatrix}$, $X = afa^{-1}$ and Y is invertible, we have

$$\ker(\overline{x}|_{\ker x}) \cap \ker(t|_{\ker x}) = 0,$$

which yields

$$\ker x \cap \ker \overline{x} \cap \ker t = 0.$$

Therefore, we have the assertion. □

Recall the fundamental isomorphism of perfect crystals (1.1). Combining the perfect crystals $B^{1,\ell}$ and $B^{n,\ell}$ with Theorem 3.2, we have

$$\Phi^1_\Lambda : \mathfrak{B}(\Lambda) \longrightarrow \mathfrak{B}(\Lambda') \otimes B^{1,\ell},$$
$$\Phi^n_\Lambda : \mathfrak{B}(\Lambda) \longrightarrow \mathfrak{B}(\Lambda'') \otimes B^{n,\ell},$$

where $\Lambda' := \sum_{i\in I} \Lambda(h_{i+1})\Lambda_i$ and $\Lambda'' := \sum_{i\in I} \Lambda(h_{i-1})\Lambda_i$. We give a geometric interpretation of the isomorphisms Φ^1_Λ and Φ^n_Λ in terms of quiver varieties.

Theorem 5.4. *Let $X \in \mathfrak{B}(\Lambda)$ and $\alpha := \Lambda - \mathrm{wt}(X)$. Set*

$$\beta = \underline{\dim}(\ker x) \quad (\textit{resp. } \gamma = \underline{\dim}(\ker \overline{x}))$$

for $[\chi = x + \overline{x}, t]$ in a nonempty open subset of X as in Lemma 5.2. Let $\phi : V(\alpha)/\ker x \to V(\alpha - \beta)$ (resp. $\bar{\phi} : V(\alpha)/\ker \overline{x} \to V(\alpha - \gamma)$) be an I-graded vector space isomorphism.

(1) *Let $\Lambda' := \sum_{i\in I} \Lambda(h_{i+1})\Lambda_i$. Then there exists a unique irreducible component $X' \in \mathfrak{B}(\Lambda')$ satisfying the following conditions:*
   (a) *there exists a nonempty open subset $U \subset X$ such that*

   $$[\phi \circ (\chi|_{V(\alpha)/\ker x}) \circ \phi^{-1},\ (t \circ x|_{V(\alpha)/\ker x}) \circ \phi^{-1}] \in X'$$

   for any element $[\chi = x + \overline{x}, t] \in U$,
   (b) *there exists a nonempty open subset $U' \subset X'$ such that every element $[\chi', t'] \in U'$ can be written as*

   $$[\chi', t'] = [\phi \circ (\chi|_{V(\alpha)/\ker x}) \circ \phi^{-1}, (t \circ x|_{V(\alpha)/\ker x}) \circ \phi^{-1}]$$

   for some element $[\chi = x + \overline{x}, t] \in X$,
   (c) *moreover, we have*

   $$\Phi^1_\Lambda(X) = X' \otimes \psi^{1,\ell}(\mathrm{wt}(b_0^\Lambda) - \mathrm{cl}(\beta)),$$

   where $\psi^{1,\ell}$ is the 1-1 map in (2.2), and $\mathbf{p}^1(\Lambda) = (b_k^\Lambda)_{k\geq 0}$ is the ground-state path of Λ in $B^{1,\ell}$ given in (2.3).

(2) *Let $\Lambda'' := \sum_{i\in I} \Lambda(h_{i-1})\Lambda_i$. Then there exists a unique irreducible component $X'' \in \mathfrak{B}(\Lambda'')$ satisfying the following conditions:*

(a) *there exists a nonempty open subset* $U \subset X$ *such that*

$$[\bar{\phi} \circ (\chi|_{V(\alpha)/\ker \overline{x}}) \circ \bar{\phi}^{-1},\ (t \circ \overline{x}|_{V(\alpha)/\ker \overline{x}}) \circ \bar{\phi}^{-1}] \in X'',$$

for any element $[\chi = x + \overline{x}, t] \in U$,

(b) *there exists a nonempty open subset* $U'' \subset X''$ *such that every element* $[\chi'', t''] \in U''$ *can be written as*

$$[\chi'', t''] = [\bar{\phi} \circ (\chi|_{V(\alpha)/\ker \overline{x}}) \circ \bar{\phi}^{-1}, (t \circ \overline{x}|_{V(\alpha)/\ker \overline{x}}) \circ \bar{\phi}^{-1}]$$

for some element $[\chi = x + \overline{x}, t] \in X$,

(c) *moreover, we have*

$$\Phi^n_\Lambda(X) = X'' \otimes \psi^{n,\ell}(\mathrm{wt}(\overline{b}^\Lambda_0) - \mathrm{cl}(\gamma)),$$

where $\psi^{n,\ell}$ *is the 1-1 map in* (2.5), *and* $\mathbf{p}^n(\Lambda) = (\overline{b}^\Lambda_k)_{k \ge 0}$ *is the ground-state path of* Λ *in* $B^{n,\ell}$ *given in* (2.6).

Note that the open sets U, U' and U'' in Theorem 5.4 do not depend on the choices of ϕ and $\bar{\phi}$ since $[\chi, t]$ denotes the $G(\alpha)$-orbit of $(\chi, t) \in X(\Lambda, \alpha)$.

Proof. We first deal with the assertion (1). Let $\mathbf{Y} = (Y^{(1)}, \ldots, Y^{(\ell)})$ be the element of $\mathcal{Y}^1(\Lambda)$ corresponding to X via the isomorphism in Proposition 4.3. Recall that $(x(\mathbf{Y}), V(\alpha))$ is the representation of (I, Ω) appeared in (4.8).

(a) From (3.3) and (4.10), there is a nonempty open subset $U \subset X$ such that, for any element $[\chi, t] \in U$, χ is contained in the conormal bundle of the $G(\alpha)$-orbit of $x(\mathbf{Y})$. For $1 \le k \le \ell$, let $Y'^{(k)}$ be the Young wall obtained from $Y^{(k)}$ by removing the 0th column $y^{(k)}_0$ of $Y^{(k)}$. It follows from Equation (4.9) that

$$\beta = \sum_{k=1}^{\ell} \mathrm{wt}(y^{(k)}_0). \tag{5.2}$$

By construction, $\mathbf{Y}' := (Y'^{(1)}, \ldots, Y'^{(\ell)})$ can be considered as the element in $\mathcal{Y}^1(\Lambda')$. Set

$$X' := \text{the irreducible component in } \mathfrak{B}(\Lambda') \text{ corresponding to } \mathbf{Y}'.$$

Take an element $[\chi, t] \in U$, and let $\chi' := \chi|_{V(\alpha)/\ker x}$ and $t' := t \circ x|_{V(\alpha)/\ker x}$. We write $\chi' = x' + \overline{x}'$. Note that χ' and t' are well-defined by Lemma 5.1. By construction, $\phi \circ x' \circ \phi^{-1}$ is contained in the $G(\alpha - \beta)$-orbit of $x(\mathbf{Y}')$. Since $x'\overline{x}' - \overline{x}'x' = (x\overline{x} - \overline{x}x)|_{V(\alpha)/\ker x} = 0$, by (3.1), we have

$$\mu_i(\phi \circ \chi' \circ \phi^{-1}) = 0 \ (i \in I).$$

Hence $[\phi \circ \chi' \circ \phi^{-1},\ t' \circ \phi^{-1}]$ is contained in X' if (χ', t') satisfies the stability condition.

Suppose that there is a nonzero χ'-invariant subspace S' of $V(\alpha)/\ker x$ with $t'(S') = 0$. Let $\pi : V(\alpha) \to V(\alpha)/\ker x$ be the natural projection. Set

$$S := x(\pi^{-1}(S')) \subset V(\alpha).$$

Since S' is nonzero, S is a nonzero I-graded subspace of $V(\alpha)$. For any element $v \in S$, v can be written as $v = xw$ for some $w \in \pi^{-1}(S')$. Note that $\chi w \in \pi^{-1}(S')$ since S' is χ'-invariant. Then, it follows from

$$\chi v = (x + \overline{x})xw = x\chi(w)$$

that S is χ-invariant. Moreover, by the assumption $t'(S') = 0$, we have $t(S) = 0$. Hence S is a nonzero χ-invariant subspace with $t(S) = 0$, which is a contradiction. Therefore, (χ', t') satisfies the stability condition.

(b) Let U' be a nonempty open subset of X' such that, for any element $[\chi', t'] \in U'$, χ' is contained in the conormal bundle of the $G(\alpha - \beta)$-orbit of $x(\mathbf{Y}')$. Take an element $[\chi' = x' + \overline{x}', t'] \in U'$. By construction, x' is contained in the $G(\alpha - \beta)$-orbit of $x(\mathbf{Y}')$. Hence, there exists an element x in the $G(\alpha)$-orbit of $x(\mathbf{Y})$ such that

$$x' = \phi \circ x|_{V(\alpha)/\ker x} \circ \phi^{-1}.$$

Then the assertion (b) follows from Lemma 5.3.

(c) Combining Proposition 4.1 with the fundamental isomorphism of perfect crystals (1.1), we have

$$\Phi : \mathcal{Y}^1(\Lambda) \longrightarrow \mathcal{Y}^1(\Lambda') \otimes B^{1,\ell}.$$

Then, by Equation (5.2) and the isomorphism (4.2), we obtained

$$\Phi(\mathbf{Y}) = \mathbf{Y}' \otimes \psi^{1,\ell}(\mathrm{wt}(b_0^{\Lambda})) - \mathrm{cl}(\beta)),$$

which yields

$$\Phi_{\Lambda}^1(X) = X' \otimes b.$$

The remaining case (2) can be obtained in the same manner. □

Let $\alpha, \beta \in Q^+$ with $\beta \leq \alpha$, and $\Lambda' = \sum_{i \in I} \Lambda(h_{i+1})\Lambda_i$. We consider an analogue of the diagram given in [20, Section 12.10]:

$$X(\Lambda', \alpha - \beta) \xleftarrow{p_1} F' \xrightarrow{p_2} F'' \xrightarrow{p_3} X(\Lambda, \alpha), \tag{5.3}$$

where F'' is the variety of all triples $(\chi = x + \overline{x}, t, M)$ such that

(a) $(\chi, t) \in X(\Lambda, \alpha)$,

(b) M is a χ-invariant subspace of $V(\alpha)$ with $\underline{\dim} M = \beta$ and $t \circ x(M) = 0$,

and F' is the variety of all quintuples (χ, t, M, f, g) such that

(a) $(\chi, t, M) \in F''$,

(b) $f = (f_i)_{i \in I}, g = (g_i)_{i \in I}$ give an exact sequence

$$0 \longrightarrow V_i(\beta) \xrightarrow{f_i} V_i(\alpha) \xrightarrow{g_i} V_i(\alpha - \beta) \longrightarrow 0$$

such that $\mathrm{Im}\, f = M$.

We also have

$$p_1(\chi, t, M, f, g) := (\tilde{g} \circ \chi|_{V(\alpha)/M} \circ \tilde{g}^{-1},\ (t \circ x|_{V(\alpha)/M}) \circ \tilde{g}^{-1}),$$

where $\chi = x + \overline{x}$, and $\tilde{g} : V(\alpha)/M \to V(\alpha - \beta)$ is the I-graded isomorphism induced by g,

$$p_2(\chi, t, M, f, g) := (\chi, t, M) \ \text{ and } \ p_3(\chi, t, M) := (\chi, t).$$

Note that p_2 is a $G(\alpha - \beta) \times G(\beta)$-principle bundle and an open map.

Take a nonempty open subset U of $X \in \mathfrak{B}(\Lambda)$ as in Lemma 5.2 and let $\alpha := \Lambda - \mathrm{wt}(X)$ and $\beta := \underline{\dim}(\ker x)$ for $[\chi = x + \overline{x}, t] \in U$. Let

$$\pi : X(\Lambda, \alpha)^{\mathrm{st}} \to X(\Lambda, \alpha)^{\mathrm{st}}/G(\alpha)$$

and $U_0 := \pi^{-1}(U)$. Note that $X(\Lambda, \alpha)^{\mathrm{st}}$ is a nonempty open subset of $X(\Lambda, \alpha)$. Define a map $\iota : U_0 \to F''$ by

$$\iota(\chi, t) = (\chi, t, \ker x)$$

for $(\chi = x + \overline{x}, t) \in U_0$. Note that $p_3 \circ \iota = \mathrm{id}|_{U_0}$. We set $X' \in \mathfrak{B}(\Lambda')$ to be the irreducible component given in Theorem 5.4 (1). Given a nonempty open subset $U_0' \subset X(\Lambda', \alpha - \beta)$ with $U_0' \cap \pi^{-1}(X') \neq \emptyset$, it follows from Lemma 5.3 that

$$\tilde{U_0} := \iota^{-1} \circ p_2 \circ p_1^{-1}(U_0')$$

is a nonempty open subset of U_0. Therefore, given a nonempty open subset $U' \subset X'$, there exists a nonempty open subset $\tilde{U} \subset X$ such that, for any element $[\chi = x + \overline{x}, t] \in \tilde{U}$,

$$[\phi \circ (\chi|_{V(\alpha)/\ker x}) \circ \phi^{-1}, (t \circ x|_{V(\alpha)/\ker x}) \circ \phi^{-1}] \in U',$$

where $\phi : V(\alpha)/\ker x \to V(\alpha - \beta)$ is an I-graded vector space isomorphism.

In the same manner, let $\Lambda'' = \sum_{i \in I} \Lambda(h_{i-1})\Lambda_i$ and $\gamma := \underline{\dim}(\ker \overline{x})$ for $[\chi = x + \overline{x}, t] \in U$. We consider the following diagram:

$$X(\Lambda'', \alpha - \gamma) \xleftarrow{\overline{p}_1} \overline{F}' \xrightarrow{\overline{p}_2} \overline{F}'' \xrightarrow{\overline{p}_3} X(\Lambda, \alpha),$$

where $\overline{F}''$ is the variety of all triples $(\chi = x + \overline{x}, t, M)$ such that

(a) $(\chi, t) \in X(\Lambda, \alpha)$,

(b) M is a χ-invariant subspace of $V(\alpha)$ with $\underline{\dim} M = \gamma$ and $t \circ \overline{x}(M) = 0$,

and $\overline{F}'$ is the variety of all quintuples (χ, t, M, f, g) such that

(a) $(\chi, t, M) \in \overline{F}''$,

(b) $f = (f_i)_{i \in I}, g = (g_i)_{i \in I}$ give an exact sequence

$$0 \longrightarrow V_i(\gamma) \xrightarrow{f_i} V_i(\alpha) \xrightarrow{g_i} V_i(\alpha - \gamma) \longrightarrow 0$$

such that $\mathrm{Im}\, f = M$.

We set

$$\overline{p}_1(\chi, t, M, f, g) := (\tilde{g} \circ \chi|_{V(\alpha)/M} \circ \tilde{g}^{-1},\ (t \circ \overline{x}|_{V(\alpha)/M}) \circ \tilde{g}^{-1}),$$

where $\chi = x + \overline{x}$, and $\tilde{g} : V(\alpha)/M \to V(\alpha - \gamma)$ is the I-graded isomorphism induced by g,

$$\overline{p}_2(\chi, t, M, f, g) := (\chi, t, M) \ \text{ and } \ \overline{p}_3(\chi, t, M) := (\chi, t).$$

Define a map $\overline{\iota} : U \to \overline{F}''$ by

$$\overline{\iota}(\chi, t) = (\chi, t, \ker \overline{x})$$

for $[\chi = x + \overline{x}, t] \in U$, and let X'' be the irreducible component given in Theorem 5.4 (2). Then we obtain that, given a nonempty open subset $U'' \subset X''$, there exists a nonempty open subset $\tilde{U} \subset X$ such that for any element $[\chi = x + \overline{x}, t] \in \tilde{U}$,

$$[\bar{\phi} \circ (\chi|_{V(\alpha)/\ker \overline{x}}) \circ \bar{\phi}^{-1}, (t \circ \overline{x}|_{V(\alpha)/\ker \overline{x}}) \circ \bar{\phi}^{-1}] \in U'',$$

where $\bar{\phi} : V(\alpha)/\ker \overline{x} \to V(\alpha - \gamma)$ is an I-graded vector space isomorphism . Consequently, we obtain the following lemma.

Lemma 5.5. *With the same notations as in Theorem 5.4, we have the following.*

(1) *Given a nonempty open subset $U' \subset X'$, there exists a nonempty open subset $\tilde{U} \subset X$ such that*

$$[\phi \circ (\chi|_{V(\alpha)/\ker x}) \circ \phi^{-1}, (t \circ x|_{V(\alpha)/\ker x}) \circ \phi^{-1}] \in U'$$

for any element $[\chi = x + \overline{x}, t] \in \tilde{U}$.

(2) *Given a nonempty open subset $U'' \subset X''$, there exists a nonempty open subset $\tilde{U} \subset X$ such that*

$$[\bar{\phi} \circ (\chi|_{V(\alpha)/\ker \overline{x}}) \circ \bar{\phi}^{-1}, (t \circ \overline{x}|_{V(\alpha)/\ker \overline{x}}) \circ \bar{\phi}^{-1}] \in U''$$

for any element $[\chi = x + \overline{x}, t] \in \tilde{U}$.

Let us recall the path realizations $\mathcal{P}^1(\Lambda)$, $\mathcal{P}^n(\Lambda)$ and $\mathcal{P}^{\mathrm{ad}}(\Lambda)$ arising from the perfect crystals $B^{1,\ell}, B^{n,\ell}$ and $B^{\mathrm{ad},\ell}$ respectively. By Theorem 1.3 and Theorem 3.2, we have crystal isomorphisms from $\mathfrak{B}(\Lambda)$ to $\mathcal{P}^1(\Lambda)$, $\mathcal{P}^n(\Lambda)$ and $\mathcal{P}^{\mathrm{ad}}(\Lambda)$ respectively. We give explicit descriptions in terms of quiver varieties.

Theorem 5.6. *Let*

$$\Upsilon^1 : \mathfrak{B}(\Lambda) \longrightarrow \mathcal{P}^1(\Lambda) \ \ (\textit{resp. } \Upsilon^n : \mathfrak{B}(\Lambda) \longrightarrow \mathcal{P}^n(\Lambda))$$

be the unique crystal isomorphism given by Theorem 1.3 and Theorem 3.2. Let $\mathbf{p}^1(\Lambda) := (b_i^\Lambda)_{i\geq 0}$ (resp. $\mathbf{p}^n(\Lambda) := (\overline{b}_i^\Lambda)_{i\geq 0}$) be the ground-state path of Λ in $B^{1,\ell}$ (resp. $B^{n,\ell}$) given in (2.3) *(resp.* (2.6)*). Take an irreducible component $X \in \mathfrak{B}(\Lambda)$. Then there exists a nonempty open subset $U \subset X$ satisfying the following conditions:*

(1) *let*

$$\mathfrak{r}_i := \mathrm{wt}(b_i^\Lambda) - \mathrm{cl}(\underline{\dim}(\ker x^{i+1}/\ker x^i)),$$

$$\mathfrak{s}_i := \mathrm{wt}(\overline{b}_i^\Lambda) - \mathrm{cl}(\underline{\dim}(\ker \overline{x}^{i+1}/\ker \overline{x}^i))$$

for $[\chi = x + \overline{x}, t] \in U$, then the weights $\mathfrak{r}_i$ and $\mathfrak{s}_i$ do not depend on the choice of $[\chi, t]$,

(2) *moreover,*

$$\Upsilon^1(X) = (\psi^{1,\ell}(\mathfrak{r}_i))_{i\geq 0} \quad \textit{and} \quad \Upsilon^n(X) = (\psi^{n,\ell}(\mathfrak{s}_i))_{i\geq 0},$$

where $\psi^{1,\ell}$ *(resp.* $\psi^{n,\ell}$*) is the 1-1 map in* (2.2) *(resp.* (2.5)*).*

Proof. Let $\alpha := \Lambda - \mathrm{wt}(X)$. We will use induction on $\mathrm{ht}(\alpha)$. If $\mathrm{ht}(\alpha) = 0$, then there is nothing to prove. Assume that $\mathrm{ht}(\alpha) > 0$. We first deal with the case of Υ^1 and $\ker x$. Let $\Lambda' := \sum_{i\in I} \Lambda(h_{i+1})\Lambda_i$, and X' be the irreducible component of $\mathfrak{B}(\Lambda')$ given in Theorem 5.4 (1). By the induction hypothesis, there is a nonempty open subset $U' \subset X'$ satisfying the conditions of our assertion. By Lemma 5.5, there is a nonempty open subset $U \subset X$ such that

$$\underline{\dim} \ker x'^{i+1} / \ker x'^{i} = \underline{\dim} \ker x^{i+2} / \ker x^{i+1}$$

for $i \in \mathbb{Z}_{>0}$, $[\chi' = x' + \overline{x}', t'] \in U'$ and $[\chi = x + \overline{x}, t] \in U$. Since the subsequence $(b_i^{\Lambda})_{i\geq 1}$ of the ground-state path $\mathbf{p}^1(\Lambda)$ can be viewed as the ground-state path of Λ' in $B^{1,\ell}$, it follows from Theorem 5.4 (1) that $\mathfrak{r}_i$ does not depend on the choice of $[\chi, t] \in U$ for each i, and

$$\Upsilon^1(X) = (\psi^{1,\ell}(\mathfrak{r}_i))_{i\geq 0}.$$

The remaining case, Υ^n and $\ker \overline{x}$, can be proved in the same manner. $\square$

Theorem 5.7. *Let*

$$\Upsilon^{\mathrm{ad}} : \mathfrak{B}(\Lambda) \longrightarrow \mathcal{P}^{\mathrm{ad}}(\Lambda)$$

be the unique crystal isomorphism given by Theorem 1.3 and Theorem 3.2. Let $\Lambda'' = \sum_{i\in I} \Lambda(h_{i-1})\Lambda_i$, *and let* $\mathbf{p}^1(\Lambda'') := (b_i^{\Lambda''})_{i\geq 0}$ *(resp.* $\mathbf{p}^n(\Lambda) := (\overline{b}_i^{\Lambda})_{i\geq 0}$*) be the ground-state path of* Λ'' *(resp.* Λ*) in* $B^{1,\ell}$ *(resp.* $B^{n,\ell}$*) given in* (2.3) *(resp.* (2.6)*). Take an irreducible component* $X \in \mathfrak{B}(\Lambda)$*. Then there exists a nonempty open subset* $U \subset X$ *satisfying the following conditions:*

(1) *let*

$$\mathfrak{r}_i := \mathrm{wt}(b_0^{\Lambda''}) - \mathrm{cl}(\underline{\dim}(\ker(x\overline{x})^{i+1} / \ker \overline{x}(x\overline{x})^i)),$$
$$\mathfrak{s}_i := \mathrm{wt}(\overline{b}_0^{\Lambda}) - \mathrm{cl}(\underline{\dim}(\ker \overline{x}(x\overline{x})^i / \ker(x\overline{x})^i))$$

for $[\chi = x + \overline{x}, t] \in U$, *then the weights* $\mathfrak{r}_i$ *and* $\mathfrak{s}_i$ *do not depend on the choice of* $[\chi, t]$,

(2) *moreover,*

$$\Upsilon^{\mathrm{ad}}(X) = (\psi^{\mathrm{ad},\ell}(\mathfrak{r}_i, \mathfrak{s}_i))_{i\geq 0},$$

where $\psi^{\mathrm{ad},\ell}$ *is the 1-1 map in* (2.8).

Proof. Let $\alpha := \Lambda - \mathrm{wt}(X)$. We will use induction on $\mathrm{ht}(\alpha)$. Since the case $\mathrm{ht}(\alpha) = 0$ is trivial, we may assume $\mathrm{ht}(\alpha) > 0$. Let $\gamma := \underline{\dim}(\ker \overline{x})$ for $[\chi = x + \overline{x}, t]$ in a nonempty open subset of X as in Lemma 5.2, and let X'' be the irreducible component of $\mathfrak{B}(\Lambda'')$ associated with X as in Theorem 5.4 (2). Similarly, let $\beta := \underline{\dim}(\ker x'')$ for $[\chi'' = x'' + \overline{x}'', t'']$ in a nonempty open subset of X'' as in Lemma 5.2, and let X' be the irreducible component of $\mathfrak{B}(\Lambda)$ associated with X'' as in Theorem 5.4 (1). Then we have

$$\Phi^1_{\Lambda''}(X'') = X' \otimes b \quad \text{and} \quad \Phi^n_{\Lambda}(X) = X'' \otimes \overline{b}$$

for some $b \in B^{1,\ell}$ and $\overline{b} \in B^{n,\ell}$. From the fundamental isomorphism (1.1) and Lemma 2.2, there is a crystal isomorphism

$$\Phi^{\mathrm{ad}}_{\Lambda} : \mathfrak{B}(\Lambda) \to \mathfrak{B}(\Lambda) \otimes B^{\mathrm{ad},\ell}.$$

By Lemma 2.1, we obtain

$$\Phi^{\mathrm{ad}}_{\Lambda}(X) = X' \otimes \xi^{\ell}(b \otimes \overline{b}).$$

On the other hand, by the induction hypothesis, there is an open subset $U' \subset X'$ satisfying the conditions of our assertion. Then it follows from Lemma 5.5 that there is a nonempty open subset $U \subset X$ such that, for $[\chi = x + \overline{x}, t] \in U$,

$$[\tilde{\phi} \circ (\chi|_{V(\alpha)/\ker x\overline{x}}) \circ \tilde{\phi}^{-1}, (t \circ x\overline{x}|_{V(\alpha)/\ker x\overline{x}}) \circ \tilde{\phi}^{-1}] \in U',$$

where $\tilde{\phi} : V(\alpha)/\ker x\overline{x} \to V(\alpha - \beta - \gamma)$ is an I-graded vector space. Note that $\gamma = \underline{\dim}(\ker \overline{x})$, $\beta = \underline{\dim}(\ker x\overline{x}/\ker \overline{x})$ and $\beta + \gamma = \underline{\dim}(\ker x\overline{x})$. By Theorem 5.4, we have

$$\mathrm{wt}(\overline{b}) = \mathrm{wt}(b_0^{\Lambda}) - \mathrm{cl}(\gamma) \quad \text{and} \quad \mathrm{wt}(b) = \mathrm{wt}(b_0^{\Lambda''}) - \mathrm{cl}(\beta),$$

which yields, by Equation (2.8),

$$\xi^{\ell}(b \otimes \overline{b}) = \psi^{\mathrm{ad},\ell}(\mathfrak{r}_0, \mathfrak{s}_0).$$

Since

$$\begin{aligned} \underline{\dim}(\ker(x\overline{x})^{i+2}/\ker \overline{x}(x\overline{x})^{i+1}) &= \underline{\dim}(\ker(x'\overline{x}')^{i+1}/\ker \overline{x}'(x'\overline{x}')^{i}), \\ \underline{\dim}(\ker \overline{x}(x\overline{x})^{i+1}/\ker \overline{x}(x\overline{x})^{i+1}) &= \underline{\dim}\overline{x}'(\ker(x'\overline{x}')^{i}/\ker \overline{x}'(x'\overline{x}')^{i}) \end{aligned}$$

for $[\chi = x + \overline{x}, t] \in U$ and $[\chi' = x' + \overline{x}', t'] \in U'$, our assertion follows from a standard induction argument. □

The following corollary, which is a consequence of Theorem 5.4 and Theorem 5.7, can be regarded as a geometric interpretation of the fundamental isomorphism of perfect crystals

$$\Phi^{\mathrm{ad}}_{\Lambda} : \mathfrak{B}(\Lambda) \xrightarrow{\sim} \mathfrak{B}(\Lambda) \otimes B^{\mathrm{ad},\ell}.$$

Corollary 5.8. *Let $X \in \mathfrak{B}(\Lambda)$ and $\alpha := \Lambda - \mathrm{wt}(X)$. Set*

$$\beta := \underline{\dim} \ker x\overline{x}/\ker \overline{x} \quad \textit{and} \quad \gamma := \underline{\dim} \ker \overline{x}$$

for $[\chi = x + \overline{x}]$ in a nonempty open subset of X as in Theorem 5.7 (1). Let $\tilde{\phi} : V(\alpha)/\ker x\overline{x} \to V(\alpha - \beta - \gamma)$ be an I-graded vector space. Then there is a unique irreducible component $X' \in \mathfrak{B}(\Lambda)$ satisfying the following conditions:

(1) *there is a nonempty open subset $U \subset X$ such that*

$$[\tilde{\phi} \circ (\chi|_{V(\alpha)/\ker x\overline{x}}) \circ \tilde{\phi}^{-1}, (t \circ x\overline{x}|_{V(\alpha)/\ker x\overline{x}}) \circ \tilde{\phi}^{-1}] \in X'$$

for any element $[\chi = x + \overline{x}, t] \in U$,

(2) *there is a nonempty open subset* $U' \subset X'$ *such that every element* $[\chi', t'] \in U'$ *can be written as*

$$[\chi', t'] = [\tilde{\phi} \circ (\chi|_{V(\alpha)/\ker x\overline{x}}) \circ \tilde{\phi}^{-1}, (t \circ x\overline{x}|_{V(\alpha)/\ker x\overline{x}}) \circ \tilde{\phi}^{-1}]$$

for some element $[\chi = x + \overline{x}, t] \in X$*,*

(3) *moreover,*

$$\Phi_{\Lambda}^{\mathrm{ad}}(X) = X' \otimes \psi^{\mathrm{ad},\ell}(\mathrm{wt}(b_0^{\Lambda''}) - \mathrm{cl}(\beta), \mathrm{wt}(\overline{b}_0^{\Lambda}) - \mathrm{cl}(\gamma)),$$

where $\psi^{\mathrm{ad},\ell}$ *is the 1-1 map in* (2.8)*, and* $\mathbf{p}^1(\Lambda'') := (b_i^{\Lambda''})_{i \geq 0}$ *(resp.* $\mathbf{p}^n(\Lambda) := (\overline{b}_i^{\Lambda})_{i \geq 0}$*) is the ground-state path of* Λ'' *(resp.* Λ*) in* $B^{1,\ell}$ *(resp.* $B^{n,\ell}$*) given in* (2.3) *(resp.* (2.6)*).*

Note that the open sets U and U' in Corollary 5.8 do not depend on the choice of $\tilde{\phi}$ since $[\chi, t]$ denotes the $G(\alpha)$-orbit of $(\chi, t) \in X(\Lambda, \alpha)$.

Example 5.9. We keep all notations in Example 4.4. By (3.3) and (4.10), it suffices to consider the fiber $\pi_{\Omega}^{-1}(x(\mathbf{Y}))$. By [20, Section 12.8, Proposition 15.5], we have

$$\begin{aligned}\pi_{\Omega}^{-1}(x(\mathbf{Y})) &= \{x(\mathbf{Y}) + \overline{x} \mid \overline{x} \in E_{\overline{\Omega}}(\alpha),\ [x(\mathbf{Y}), \overline{x}] = 0\} \\ &= \{x(\mathbf{Y}) + \overline{x} \mid \overline{x} = \overline{x}(a_1, \cdots, a_{29}),\ a_1, \ldots, a_{29} \in \mathbb{C}\}.\end{aligned}$$

Here,

$$\overline{x}(a_1, \cdots, a_{29}) := \begin{pmatrix} 0 & \overline{x}_1 & 0 \\ 0 & 0 & \overline{x}_2 \\ \overline{x}_0 & 0 & 0 \end{pmatrix} \in E_{\overline{\Omega}}(\alpha),$$

$$\overline{x}_0 := \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & a_{25} \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \overline{x}_1 := \begin{pmatrix} a_1 & a_2 & a_3 & a_4 & 0 & a_5 & a_6 \\ a_7 & a_8 & a_9 & a_{10} & 0 & a_{11} & a_{12} \\ a_{26} & 0 & a_{27} & 0 & 0 & a_{28} & a_{29} \\ a_{13} & a_{14} & a_{15} & a_{16} & 0 & a_{17} & a_{18} \end{pmatrix},$$

$$\overline{x}_2 := \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a_{19} & 0 & 0 & a_{20} \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a_{21} & 0 & 0 & a_{22} \\ a_{26} & a_{27} & a_{23} & a_{28} & a_{29} & a_{24} \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & a_{25} \end{pmatrix}$$

for $a_1, \ldots, a_{29} \in \mathbb{C}$. Let

$$x := x(\mathbf{Y}), \quad \overline{x} := \overline{x}(a_1, \cdots, a_{29})$$

and consider $a_1, \dots, a_{29}$ as indeterminates. Then we have

$$\underline{\dim}(\ker x^k) = \begin{cases} 0 & \text{if } k=0, \\ 3\alpha_0+3\alpha_1+2\alpha_2 & \text{if } k=1, \\ 4\alpha_0+4\alpha_1+5\alpha_2 & \text{if } k=2, \\ 4\alpha_0+6\alpha_1+6\alpha_2 & \text{if } k=3, \\ 4\alpha_0+7\alpha_1+6\alpha_2 & \text{otherwise,} \end{cases} \quad \text{and} \quad \underline{\dim}(\ker \overline{x}^k) = \begin{cases} 0 & \text{if } k=0, \\ 3\alpha_0+3\alpha_1+3\alpha_2 & \text{if } k=1, \\ 3\alpha_0+6\alpha_1+4\alpha_2 & \text{if } k=2, \\ 4\alpha_0+6\alpha_1+5\alpha_2 & \text{if } k=3, \\ 4\alpha_0+7\alpha_1+5\alpha_2 & \text{if } k=4, \\ 4\alpha_0+7\alpha_1+6\alpha_2 & \text{otherwise.} \end{cases}$$

Hence, by Theorem 5.6, we obtain

$$\begin{aligned} \Upsilon^1(X) &= \cdots \otimes \begin{array}{|c|c|c|}\hline 2&2&3\\\hline\end{array} \otimes \begin{array}{|c|c|c|}\hline 2&3&3\\\hline\end{array} \otimes \begin{array}{|c|c|c|}\hline 2&2&3\\\hline\end{array} \otimes \begin{array}{|c|c|c|}\hline 3&3&3\\\hline\end{array} \otimes \begin{array}{|c|c|c|}\hline 1&2&3\\\hline\end{array}, \\ \Upsilon^n(X) &= \cdots \otimes \begin{array}{|c|c|c|}\hline \overline{3}&\overline{3}&\overline{1}\\\hline\end{array} \otimes \begin{array}{|c|c|c|}\hline \overline{2}&\overline{2}&\overline{2}\\\hline\end{array} \otimes \begin{array}{|c|c|c|}\hline \overline{1}&\overline{1}&\overline{1}\\\hline\end{array} \otimes \begin{array}{|c|c|c|}\hline \overline{3}&\overline{3}&\overline{2}\\\hline\end{array} \otimes \begin{array}{|c|c|c|}\hline \overline{1}&\overline{1}&\overline{1}\\\hline\end{array} \otimes \begin{array}{|c|c|c|}\hline \overline{2}&\overline{1}&\overline{1}\\\hline\end{array}. \end{aligned}$$

On the other hand, the ground-state path of $\Lambda'' = 2\Lambda_1 + \Lambda_2$ in $B^{1,\ell}$ is given as follows.

$$\cdots \otimes \begin{array}{|c|c|c|}\hline 1&1&2\\\hline\end{array} \otimes \begin{array}{|c|c|c|}\hline 2&2&3\\\hline\end{array} \otimes \begin{array}{|c|c|c|}\hline 1&3&3\\\hline\end{array} \otimes \begin{array}{|c|c|c|}\hline 1&1&2\\\hline\end{array}.$$

By a direct computation, we have

$$\underline{\dim}(\ker (x\overline{x})^k) = \begin{cases} 0 & \text{if } k=0, \\ 4\alpha_0+6\alpha_1+5\alpha_2 & \text{if } k=1, \\ 4\alpha_0+7\alpha_1+6\alpha_2 & \text{otherwise,} \end{cases}$$

$$\underline{\dim}(\ker \overline{x}(x\overline{x})^k) = \begin{cases} 3\alpha_0+3\alpha_1+3\alpha_2 & \text{if } k=0, \\ 4\alpha_0+7\alpha_1+5\alpha_2 & \text{if } k=1, \\ 4\alpha_0+7\alpha_1+6\alpha_2 & \text{otherwise.} \end{cases}$$

By Theorem 5.7, we obtain

$$\Upsilon^{\mathrm{ad}}(X) = \cdots \otimes \begin{array}{|c|c|}\hline 1&2\\\hline 3&\multicolumn{1}{c}{}\\\cline{1-1}\end{array} \otimes \begin{array}{|c|c|}\hline 1&2\\\hline 3&\multicolumn{1}{c}{}\\\cline{1-1}\end{array} \otimes \begin{array}{|c|c|}\hline 2&3\\\hline 3&\multicolumn{1}{c}{}\\\cline{1-1}\end{array} \otimes \begin{array}{|c|c|c|c|c|c|}\hline 1&2&2&2&2&3\\\hline 3&3&3&\multicolumn{3}{c}{}\\\cline{1-3}\end{array}.$$

References

1. G. Benkart, I. Frenkel, S.-J. Kang, and H. Lee, *Level 1 perfect crystals and path realizations of basic representations at* $q = 0$, Int. Math. Res. Not. (2006), 1–28.
2. E. Date, M. Jimbo, A. Kuniba, T. Miwa, and M. Okado, *Paths, Maya diagrams and representations of* $\widehat{sl}(r, \mathbb{C})$, Adv. Stud. Pure Math. **19** (1989), 149–191.
3. O. Foda, B. Leclerc, M. Okado, J.-Y. Thibon, and T. A. Welsh, *Branching functions of* $A^{(1)}_{n-1}$ *and Jantzen-Seitz problem for Ariki-Koike algebras*, Adv. Math. **141** (1999), no. 2, 322–365.
4. I. B. Frenkel and A. Savage, *Bases of representations of type A affine Lie algebras via quiver varieties and statistical mechanics*, Int. Math. Res. Not. (2003), no. 28, 1521–1547.
5. J. Hong and S.-J. Kang, *Introduction to Quantum Groups and Crystal Bases*, Grad. Stud. Math., 42, American Mathematical Society, Providence, RI, 2002.
6. N. Jacon and C. Lecouvey, *Kashiwara and Zelevinsky involutions in affine type A*, Pacific J. Math. **243** (2009), no. 2, 287–311.

7. M. Jimbo, K. C. Misra, T. Miwa, and M. Okado, *Combinatorics of representations of $U_q(\widehat{sl}(n))$ at $q = 0$*, Comm. Math. Phys. **136** (1991), 543–566.
8. S.-J. Kang, *Crystal bases for quantum affine algebras and combinatorics of Young walls*, Proc. London Math. Soc. (3) **86** (2003), no. 1, 29–69.
9. S.-J. Kang, M. Kashiwara, and K. C. Misra, *Crystal bases of Verma modules for quantum affine Lie algebras*, Compositio Math. **92** (1994), no. 3, 299–325.
10. S.-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima, and A. Nakayashiki, *Affine crystals and vertex models*, Infinite Analysis, Part A,B(Kyoto, 1991), 449–484, Adv. Ser. Math. Phys. **16**, World Sci. Publ., River Edge, NJ (1992).
11. ______, *Perfect crystals of quantum affine Lie algebras*, Duke Math. J. **68** (1992), 499–607.
12. S.-J. Kang and H. Lee, *Higher level affine crystals and Young walls*, Algebr. Represent. Theory **9** (2006), no. 6, 593–632.
13. S.-J. Kang and E. Park, *Quiver varieties and path realizations arising from adjoint crystals of type $A_n^{(1)}$*, Trans. Amer. Math. Soc. **363** (2011), no. 10, 5341–5366.
14. M. Kashiwara, *Crystalizing the q-analogue of universal enveloping algebras*, Comm. Math. Phys. **133** (1990), no. 2, 249–260.
15. ______, *On crystal bases of the q-analogue of universal enveloping algebras*, Duke Math. J. **63** (1991), no. 2, 465–516.
16. M. Kashiwara and T. Nakashima, *Crystal graphs for representations of the q-analogue of classical Lie algebras*, J. Algebra **165** (1994), no. 2, 295–345.
17. M. Kashiwara and Y. Saito, *Geometric construction of crystal bases*, Duke Math. J. **89** (1997), no. 1, 9–36.
18. R. Kodera, *A generalization of adjoint crystals for the quantized affine algebras of type $A_n^{(1)}, C_n^{(1)}$ and $D_{n+1}^{(2)}$*, J. Algebraic Combin. **30** (2009), no. 4, 491–514.
19. G. Lusztig, *Canonical bases arising from quantized enveloping algebras*, J. Amer. Math. Soc. **3** (1990), no. 2, 447–498.
20. ______, *Quivers, perverse sheaves, and quantized enveloping algebras*, J. Amer. Math. Soc. **4** (1991), no. 2, 365–421.
21. K. C. Misra and T. Miwa, *Crystal base for the basic representation of $U_q(sl(n))$*, Comm. Math. Phys. **134** (1990), no. 1, 79–88.
22. H. Nakajima, *Instantons on ALE spaces, quiver varieties, and Kac-Moody algebras*, Duke Math. J. **76** (1994), no. 2, 365–416.
23. ______, *Quiver varieties and Kac-Moody algebras*, Duke Math. J. **91** (1998), no. 3, 515–560.
24. Y. Saito, *Crystal bases and quiver varieties*, Math. Ann. **324** (2002), no. 4, 675–688.
25. A. Savage, *Geometric and combinatorial realizations of crystal graphs*, Algebr. Represent. Theory **9** (2006), no. 2, 161–199.
26. A. Schilling and P. Sternberg, *Finite-dimensional crystals $B^{2,s}$ for quantum affine algebras of type $D_n^{(1)}$*, J. Algebraic Combin. **23** (2006), no. 4, 317–354.

School of Mathematics, Korea Institute for Advanced Study, 87 Hoegiro, Dongdaemun-gu, Seoul 130-722, Korea

E-mail address: `eypark@kias.re.kr`